arXiv:1410.5717v1 [math.AP] 21 Oct 2014

Semilinear pseudodifferential equations in spaces of tempered ultradistributions

Marco Cappiello [a] *, Stevan Pilipović [b] and Bojan Prangoski [c]

Abstract

We study a class of semilinear elliptic equations on spaces of tempered ultradistributions of Beurling and Roumieu type. Assuming that the linear part of the equation is an elliptic pseudodifferential operator of infinite order with a sub-exponential growth of its symbol and that the non linear part is given by an infinite sum of powers of u with sub-exponential growth with respect to u, we prove a regularity result in the functional setting of the quoted ultradistribution spaces for a weak Sobolev type solution u.

0 Introduction

In this paper we consider a class of semilinear equations and prove a result of regularity in the spaces of tempered ultradistributions of Beurling and Roumieu type. These ultradistributions can be regarded as a global counterpart on $\mathbb{R}^d$ of the local ultradistributions studied by Komatsu [14, 15, 17] and they represent a natural generalisation of non-quasi-analytic Gelfand-Shilov type ultradistributions, cf. [13, 19, 20]. As well as the Gelfand-Shilov spaces, they are also a good functional setting for global pseudodifferential operators of infinite order, namely with symbol $a(x,\xi)$ admitting sub-exponential growth in both x and ξ, see [2, 3, 23]. Here we want to apply the pseudodifferential operators introduced by the third author in [23] to the study of semilinear equations of the form

$$Au = f + F[u] \tag{0.1}$$

where $A = a(x,D)$, f is a given test function in our setting and $F[u]$ is a nonlinear term given by a suitable infinite sum of powers of u. In [21] we investigated the class of operators of [23] in the context of the Weyl and the Anti Wick calculus while in the recent paper [8], we considered the case of linear equations and proved a result of hypoellipticity via the construction of a parametrix. To treat semilinear equations, we need to adopt a more sophisticated method based on suitable commutators and nonlinear estimates. This method, previously used in [1] [4]-[7], [11] in the case of symbols corresponding to differential operators and of nonlinear terms of finite

2010 *Mathematical Subject Classification: 47G30, 46F05, 35A17*
keywords: Tempered ultradistributions, pseudodifferential operators, semilinear equations
* Corresponding author: marco.cappiello@unito.it

order, is used also here but with a more advanced technique since we consider infinite series both in the case of symbols and in the case of nonlinear terms. Examples of our elliptic symbols are $a(x,\xi) = e^{c\langle(x,\xi)\rangle^{1/m}}, m > 1, c \in \mathbb{R}$, whereas for what concerns the nonlinear terms we can consider $F[u] = \sum_\beta c_\beta P_\beta(x) u^{|\beta|}$ where $P_\beta(x)$ are ultrapolynomials of the form $\sum_\gamma \tilde{c}_\gamma x^\gamma/\gamma!^m, m > 1$ and c_β are suitable complex numbers tending rapidly to zero. Actually, we will consider nonlinear terms also with the additional assumption that $u \in H^s(\mathbb{R}^d), s > d/2$. With this we have plenty of examples, for example $F[u] = P(x)\cos u$ or $F[u] = P(x)e^{u^k}, k \in \mathbb{Z}_+$, where P has sub-exponential growth of the order related to the order of the growth of the symbol. In this way we can analyse elliptic operators A of infinite order and sub-exponential growth as well as nonlinear terms with sub-exponential growth, not considered in the literature, which shows an intrinsic connection of the pseudodifferential calculus of [23] with the spaces of ultradistributions.

The paper is organised as follows. In the next Section 1 we introduce the main tools involved in the paper and state the main result, namely Theorem 1.2. Section 2 contains some examples of elliptic operators and nonlinear terms which motivate our analysis and for which Theorem 1.2 holds. In Section 3 we refine some results about the pseudodifferential operators studied in [23] and we prove some precise estimates for the norms of some composed operators which will be instrumental in the proof of Theorem 1.2. Finally, Section 4 is devoted to the proof of the theorem which will be divided in two parts, one corresponding to the proof of the decay properties of the solution and the other related to its regularity.

1 Notation and the main theorem

Before stating our results, let us fix some notation and introduce the functional setting where they are obtained. In the sequel, the sets of integer, non-negative integer, positive integer, real and complex numbers are denoted as standard by $\mathbb{Z}$, $\mathbb{N}$, $\mathbb{Z}_+$, $\mathbb{R}$, $\mathbb{C}$. We denote $\langle x\rangle = (1+|x|^2)^{1/2}$ for $x \in \mathbb{R}^d$, $D^\alpha = D_1^{\alpha_1}\dots D_d^{\alpha_d}$, $D_j^{\alpha_j} = i^{-1}\partial^{\alpha_j}/\partial x^{\alpha_j}$, $\alpha = (\alpha_1,\dots,\alpha_d) \in \mathbb{N}^d$. Fixed $B > 0$, we shall denote by Q_B^c the set of all $(x,\xi) \in \mathbb{R}^{2d}$ for which we have $\langle x\rangle \geq B$ or $\langle\xi\rangle \geq B$. Finally, for $s \in \mathbb{R}$, we shall denote by $H^s(\mathbb{R}^d)$ the Sobolev space of all $u \in \mathcal{S}'(\mathbb{R}^d)$ for which $\langle\xi\rangle^s \hat{u}(\xi) \in L^2(\mathbb{R}^d)$, where $\hat{u}$ denotes the Fourier transform of u. Following [14], in the sequel we shall consider sequences M_p of positive numbers such that $M_0 = M_1 = 1$ and satisfying all or some of the following conditions: $(M.1)$ $M_p^2 \leq M_{p-1}M_{p+1}$, $p \in \mathbb{Z}_+$; $(M.2)$ $M_p \leq c_0 H^p \min_{0\leq q\leq p}\{M_{p-q}M_q\}$, $p,q \in \mathbb{N}$, for some $c_0, H \geq 1$; $(M.3)$ $\sum_{p=q+1}^{\infty} M_{p-1}/M_p \leq c_0 q M_q/M_{q+1}$, $q \in \mathbb{Z}_+$; $(M.4)$ $(M_p/p!)^2 \leq M_{p-1}/(p-1)!\cdot M_{p+1}/(p+1)!$, for all $p \in \mathbb{Z}_+$. In some assertions in the sequel we could replace $(M.3)$ by the weaker assumption: $(M.3)'$ $\sum_{p=1}^{\infty} M_{p-1}/M_p < \infty$ (cf. [14]). We observe moreover that $(M.4)$ implies $(M.1)$. As an example of sequence satisfying all the conditions above we can take $M_p = p!^s$, $s > 1$. For a multi-index $\alpha \in \mathbb{N}^d$, M_α will mean $M_{|\alpha|}$, $|\alpha| = \alpha_1 + \dots + \alpha_d$. We can associate to any sequence M_p as above the function

$M(\rho) = \sup_{p\in\mathbb{N}} \log_+ \rho^p/M_p, \quad \rho > 0$. This is a non-negative, continuous, monotonically increasing function which vanishes for sufficiently small $\rho > 0$ and increases more rapidly than $\ln \rho^p$ when ρ tends to infinity, for any $p \in \mathbb{N}$ (cf. [14]).

As in [14], see also [20], we shall denote by $\mathfrak{R}$ the set of positive sequences which monotonically increase to infinity. For $(r_p) \in \mathfrak{R}$, consider the sequence $N_0 = 1$, $N_p = M_p R_p$, $p \in \mathbb{Z}_+$, where we denote $R_p = \prod_{j=1}^p r_j$ (in the future we will often use this notation). It is easy to verify that this sequence satisfies $(M.1)$ and $(M.3)'$. Its associated function will be denoted by $N_{r_p}(\rho)$, i.e. $N_{r_p}(\rho) = \sup_{p\in\mathbb{N}} \log_+ \rho^p/(M_p R_p)$, $\rho > 0$. Note, for given (r_p) and every $k > 0$ there is $\rho_0 > 0$ such that $N_{r_p}(\rho) \leq M(k\rho)$ for $\rho > \rho_0$.

Now we can introduce the space of tempered ultradistributions and its test function space. For $m > 0$ and a sequence M_p satisfying the conditions $(M.1) - (M.3)$, we shall denote by $\mathcal{S}_\infty^{M_p,m}(\mathbb{R}^d)$ the Banach space of all functions $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ such that

$$\|\varphi\|_m := \sup_{\alpha\in\mathbb{N}^d} \sup_{x\in\mathbb{R}^d} \frac{m^{|\alpha|}|D^\alpha\varphi(x)|e^{M(m|x|)}}{M_\alpha} < \infty, \tag{1.1}$$

endowed with the norm in (1.1) and we denote $\mathcal{S}^{(M_p)}(\mathbb{R}^d) = \varprojlim_{m\to\infty} \mathcal{S}_\infty^{M_p,m}(\mathbb{R}^d)$ and $\mathcal{S}^{\{M_p\}}(\mathbb{R}^d) = \varinjlim_{m\to 0} \mathcal{S}_\infty^{M_p,m}(\mathbb{R}^d)$. In the sequel we shall consider simultaneously the two latter spaces by using the common notation $\mathcal{S}^*(\mathbb{R}^d)$. For each space we will consider a suitable symbol class. Definitions and statements will be formulated first for the (M_p) case and then for the $\{M_p\}$ case, using the notation $*$. We shall denote by $\mathcal{S}^{*\prime}(\mathbb{R}^d)$ the strong dual space of $\mathcal{S}^*(\mathbb{R}^d)$. We refer to [12, 19, 20, 22] for the properties of $\mathcal{S}^*(\mathbb{R}^d)$ and $\mathcal{S}^{*\prime}(\mathbb{R}^d)$. Here we just recall that the Fourier transformation is an automorphism on $\mathcal{S}^*(\mathbb{R}^d)$ and on $\mathcal{S}^{*\prime}(\mathbb{R}^d)$ and that for $M_p = p!^s$, $s > 1$, we have $M(\rho) \sim \rho^{1/s}$. In this case $\mathcal{S}^*(\mathbb{R}^d)$ coincides respectively with the Gelfand-Shilov spaces $\Sigma_s(\mathbb{R}^d)$ (resp. $\mathcal{S}_s(\mathbb{R}^d)$) of all functions $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ such that

$$\sup_{\alpha,\beta\in\mathbb{N}^d} h^{-|\alpha|-|\beta|}(\alpha!\beta!)^{-s} \sup_{x\in\mathbb{R}^d} |x^\beta \partial^\alpha \varphi(x)| < \infty$$

for every $h > 0$ (resp. for some $h > 0$), cf. [13, 19]. A measurable function f on $\mathbb{R}^d$ is said to be of ultrapolynomial growth of class * if $\|f(\cdot)e^{-M(m|\cdot|)}\|_{L^\infty(\mathbb{R}^d)} < \infty$ for some $m > 0$ (resp. for every $m > 0$).

Following [23] we now introduce the class of pseudodifferential operators involved in the sequel. Let M_p, A_p be two sequences of positive numbers. We assume that M_p satisfies $(M.1)$, $(M.2)$, $(M.3)$ and $(M.4)$ and that A_p satisfies $A_0 = A_1 = 1$, $(M.1), (M.2), (M.3)'$ and $(M.4)$. Moreover we suppose that $A_p \subset M_p$ i.e. there exist $c_0 > 0, L > 0$ such that $A_p \leq c_0 L^p M_p$ for all $p \in \mathbb{N}$. Let $\rho_0 = \inf\{\rho \in \mathbb{R}_+ |\, A_p \subset M_p^\rho\}$. Obviously $0 < \rho_0 \leq 1$. Let $\rho \in \mathbb{R}_+$ be arbitrary but fixed such that $\rho_0 \leq \rho \leq 1$ if the infimum can be attained, or otherwise $\rho_0 < \rho \leq 1$. For any fixed $h > 0, m > 0$ we denote by $\Gamma_{A_p,\rho}^{M_p,\infty}(\mathbb{R}^{2d}; h, m)$ the space of all functions $a(x,\xi) \in \mathcal{C}^\infty(\mathbb{R}^{2d})$ such that

$$\sup_{\alpha,\beta\in\mathbb{Z}_+^d} \sup_{(x,\xi)\in\mathbb{R}^{2d}} \frac{|D_\xi^\alpha D_x^\beta a(x,\xi)|\langle(x,\xi)\rangle^{\rho|\alpha+\beta|} e^{-M(m|x|)-M(m|\xi|)}}{h^{|\alpha+\beta|}A_\alpha A_\beta} < \infty,$$

where $M(\cdot)$ is the associated function for the sequence M_p. Then we define

$$\Gamma^{(M_p),\infty}_{A_p,\rho}(\mathbb{R}^{2d};m)=\varinjlim_{h\to 0}\Gamma^{M_p,\infty}_{A_p,\rho}(\mathbb{R}^{2d};h,m);\quad \Gamma^{(M_p),\infty}_{A_p,\rho}(\mathbb{R}^{2d})=\varinjlim_{m\to\infty}\Gamma^{(M_p),\infty}_{A_p,\rho}(\mathbb{R}^{2d};m);$$

$$\Gamma^{\{M_p\},\infty}_{A_p,\rho}(\mathbb{R}^{2d};h)=\varinjlim_{m\to 0}\Gamma^{M_p,\infty}_{A_p,\rho}(\mathbb{R}^{2d};h,m);\quad \Gamma^{\{M_p\},\infty}_{A_p,\rho}(\mathbb{R}^{2d})=\varinjlim_{h\to\infty}\Gamma^{\{M_p\},\infty}_{A_p,\rho}(\mathbb{R}^{2d};h).$$

We associate to any symbol $a\in\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ a pseudodifferential operator $a(x,D)$ acting continuously on $\mathcal{S}^*(\mathbb{R}^d)$ and on $\mathcal{S}^{*\prime}(\mathbb{R}^d)$. A symbolic calculus for $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ (denoted there by $\Gamma^{*,\infty}_{A_p,A_p,\rho}(\mathbb{R}^{2d})$) has been constructed in [23]. As a consequence it was proved that the class of pseudodifferential operators with symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ is closed with respect to composition and adjoints, cf. [23] and the next section for details. Moreover, in [8] we consider hypoelliptic symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ and we proved the existence of parametrices for the associated operators. Now we need to introduce a notion of elliptic symbol in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$. For this purpose let $\tilde{M}_p$ be another sequence such that $\tilde{M}_0=\tilde{M}_1=1$ and satisfying $(M.1)$, $(M.2)$, $(M.3)'$ and $(M.4)$ and $M_p\subset\tilde{M}_p$, i.e. there exists $\tilde{c},\tilde{L}>0$ such that $M_p\le\tilde{c}\tilde{L}^p\tilde{M}_p$ (observe that $\tilde{M}_p$ can be the same as M_p). Obviously, without losing generality, we can assume that the constant H from $(M.2)$ is the same for the sequences A_p, M_p and $\tilde{M}_p$. For $(k_p)\in\mathfrak{R}$ we denote by $\tilde{N}_{k_p}(\cdot)$ the associated function to the sequence $\tilde{M}_p\prod_{j=1}^p k_j$. One easily obtains the following inequalities

$$\tilde{M}(\lambda/\tilde{L})\le M(\lambda)+\ln_+\tilde{c}\text{ and }\tilde{N}_{k_p}(\lambda/\tilde{L})\le N_{k_p}(\lambda)+\ln_+\tilde{c},\ \forall\lambda>0. \tag{1.2}$$

If $(k_p)\in\mathfrak{R}$ satisfies the condition $K_{p+q}\le cH^{p+q}K_pK_q$ for some $c,H\ge 1$, where K_p stands for $\prod_{j=1}^p k_j$ then the sequences M_pK_p and $\tilde{M}_pK_p$ satisfy $(M.2)$ (since M_p and $\tilde{M}_p$ do). Lemma 2.3 of [22] proves that there are plenty of sequences of this type (in fact, the quoted lemma claims that given a sequence of $\mathfrak{R}$ one can find such sequence which is smaller than the chosen one). Because of this property we will say that the sequence (k_p) satisfies $(M.2)$ (although the precise statement will be to say that the sequence $K_p=\prod_{j=1}^p k_j$ satisfies $(M.2)$).

Definition 1.1. *Given $A_p,M_p,\tilde{M}_p$ as before, a symbol $a\in\Gamma^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ is said to be $(\tilde{M}_p)$-elliptic, (resp. $\{\tilde{M}_p\}$-elliptic) if*

i) there exist $m,B,c>0$ (resp. there exist $(k_p)\in\mathfrak{R}$ which satisfies $(M.2)$ and $B,c>0$) such that

$$|a(x,\xi)|\ge ce^{\tilde{M}(m|\xi|)}e^{\tilde{M}(m|x|)}(\text{ resp. }|a(x,\xi)|\ge ce^{\tilde{N}_{k_p}(|\xi|)}e^{\tilde{N}_{k_p}(|x|)}),\quad (x,\xi)\in Q^c_B;$$

ii) for every $h>0$ there exists $C>0$ (resp. there exist $h,C>0$) such that

$$\left|D^\alpha_\xi D^\beta_x a(x,\xi)\right|\le C\frac{h^{|\alpha|+|\beta|}A_{\alpha+\beta}|a(x,\xi)|}{\langle(x,\xi)\rangle^{\rho(|\alpha|+|\beta|)}},\quad (x,\xi)\in Q^c_B.$$

Given $(k_p) \in \mathfrak{R}$ and $\gamma \in \mathbb{N}^d$, from now on we will always use the notation K_p for $\prod_{j=1}^{p} k_j$ and K_γ for $K_{|\gamma|}$.

Finally we introduce the class of nonlinear terms involved in our equations.

For $\beta \in \mathbb{N}^d$, let $p_\beta(x)$ be smooth functions on $\mathbb{R}^d$ such that for every $h > 0$ there exists $C > 0$ such that

$$|D_x^\alpha p_\beta(x)| \leq C \frac{h^{|\alpha|+|\beta|} A_\alpha e^{\tilde{M}(h|x|)}}{\tilde{M}_\alpha} \text{ for all } \alpha, \beta \in \mathbb{N}^d, \tag{1.3}$$

(respectively

$$|D_x^\alpha p_\beta(x)| \leq C \frac{h^{|\alpha|+|\beta|} A_\alpha e^{\tilde{N}_{k_p}(h|x|)}}{\tilde{M}_\alpha \prod_{j=1}^{|\alpha|} k_j} \text{ for all } \alpha, \beta \in \mathbb{N}^d.) \tag{1.4}$$

For such a family of functions $p_\beta(x)$ and $u \in H^s(\mathbb{R}^d), s > d/2$ we can consider the function

$$F[u] = \sum_{|\beta|=2}^{\infty} p_\beta u^{|\beta|}, \tag{1.5}$$

The condition $s > d/2$ implies that $F[u]$ is well defined and continuous on $\mathbb{R}^d$ and $\left\| F[u] e^{-\tilde{M}(h|\cdot|)} \right\|_{L^\infty(\mathbb{R}^d)} < \infty$ (resp. $\left\| F[u] e^{-\tilde{N}_{k_p}(h|\cdot|)} \right\|_{L^\infty(\mathbb{R}^d)} < \infty$) for some h. This (together with (1.2)) implies that $F[u] \in \mathcal{S}^{*\prime}(\mathbb{R}^d)$.

The main result of the paper is the following one.

Theorem 1.2. *Let $a \in \Gamma^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) and let $f \in \mathcal{S}^*\left(\mathbb{R}^d\right)$. Let $u \in H^s\left(\mathbb{R}^d\right)$, $s > d/2$, be a solution of the equation* (0.1) *with $F[u]$ defined by* (1.3) *and* (1.5) *(resp.* (1.4) *and* (1.5)*). Then the following properties hold:*

i) For every $h > 0$ there exists $C > 0$ (resp. there exist $h, C > 0$) such that $|u(x)| \leq Ce^{-M(h|x|)}$. Moreover, $u \in \mathcal{C}^\infty(\mathbb{R}^d)$ with the following estimate on its derivatives: there exists $\tilde{h} > 0$ such that

$$\sup_\alpha \frac{\tilde{h}^{|\alpha|} \|D^\alpha u\|_{L^\infty}}{\tilde{M}_\alpha} < \infty, \left(\text{ resp. } \sup_\alpha \frac{\tilde{h}^{|\alpha|} \|D^\alpha u\|_{L^\infty}}{\tilde{M}_\alpha \prod_{j=1}^{|\alpha|} k_j} < \infty \right).$$

ii) Furthermore, if $F[u]$ is a finite sum, then $u \in \mathcal{S}^*\left(\mathbb{R}^d\right)$.

2 Examples

In this section we give an example of a non standard elliptic operator a, given by (2.1) below, and of nonlinear terms to which Theorem 1.2 can be applied when $M_p = p!^l$, $l > 1$. This examples underline the novelties of this paper with respect to the results obtained in [4, 5, 6].

Example 1. Let $A_p = p!^v$, with $1 < v < l$ and $0 < \rho < 1$ is such that $v \leq l\rho$. Let $a_0 : (0,\infty) \to (0,\infty)$ be given by $a_0(\lambda) = \sum_{n=0}^{\infty} \frac{h^n \lambda^n}{n!^{l+l'}}$. Then

$$\begin{aligned} a_0^{(k)}(\lambda) &= \frac{1}{\lambda^k} \sum_{n\geq k} \frac{n!}{(n-k)!} \cdot \frac{h^n \lambda^n}{n!^{l+l'}} \\ &= \frac{h^{k/(l+l')}}{\lambda^{k(l+l'-1)/(l+l')}} \sum_{n\geq k} \left(\frac{h^n \lambda^n}{n!^{l+l'}}\right)^{(l+l'-1)/(l+l')} \cdot \left(\frac{h^{n-k}\lambda^{n-k}}{(n-k)!^{l+l'}}\right)^{1/(l+l')} \\ &\leq \frac{h^{k/(l+l')}}{\lambda^{k(l+l'-1)/(l+l')}} \left(\sum_{n\geq k} \frac{h^n \lambda^n}{n!^{l+l'}}\right)^{(l+l'-1)/(l+l')} \cdot \left(\sum_{n\geq k} \frac{h^{n-k}\lambda^{n-k}}{(n-k)!^{l+l'}}\right)^{1/(l+l')} \\ &\leq \frac{h^{k/(l+l')}}{\lambda^{k(l+l'-1)/(l+l')}} a_0(\lambda), \end{aligned}$$

where, in the first inequality we used Holder's inequality with $p = (l+l')/(l+l'-1)$ and $q = l + l'$. Define

$$a(w) = a_0(\langle w \rangle),\ w \in \mathbb{R}^{2d} \tag{2.1}$$

Then $a \in \mathcal{C}^\infty(\mathbb{R}^{2d})$. To prove that $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ and that it is elliptic, we need some preliminary results. First we recall the following multidimensional variant of the Faà di Bruno formula (see [10, Corollary 2.10]).

Proposition 2.1. *([10]) Let $|\alpha| = n \geq 1$ and $h(x_1,...,x_d) = f(g(x_1,...,x_d))$ with $g \in \mathcal{C}^n$ in a neighbourhood of x^0 and $f \in \mathcal{C}^n$ in a neighbourhood of $y^0 = g(x^0)$. Then*

$$\partial^\alpha h(x^0) = \sum_{r=1}^{n} f^{(r)}(y^0) \sum_{p(\alpha,r)} \alpha! \prod_{j=1}^{n} \frac{\left(\partial^{\alpha^{(j)}} g(x^0)\right)^{k_j}}{k_j! \left(\alpha^{(j)}!\right)^{k_j}},$$

where

$$\begin{aligned} p(\alpha,r) = \Bigg\{ & \left(k_1,...,k_n;\alpha^{(1)},...,\alpha^{(n)}\right) \Big| \text{ for some } 1 \leq s \leq n, k_j = 0 \text{ and } \alpha^{(j)} = 0 \\ & \text{for } 1 \leq j \leq n-s;\ k_j > 0 \text{ for } n-s+1 \leq j \leq n; \text{ and} \\ & 0 \prec \alpha^{(n-s+1)} \prec ... \prec \alpha^{(n)} \text{ are such that} \\ & \sum_{j=1}^{n} k_j = r,\ \sum_{j=1}^{n} k_j \alpha^{(j)} = \alpha \Bigg\}. \end{aligned}$$

The relation $\prec$ used in this proposition is defined in the following way (cf. [10]). We say that $\beta \prec \alpha$ when one of the following holds:

(*i*) $|\beta| < |\alpha|$;

(*ii*) $|\beta| = |\alpha|$ and $\beta_1 < \alpha_1$;

(*iii*) $|\beta| = |\alpha|$, $\beta_1 = \alpha_1, ..., \beta_k = \alpha_k$ and $\beta_{k+1} < \alpha_{k+1}$ for some $1 \leq k < d$.

Before we continue, we need the following technical result.

Lemma 2.2. *For $\beta \in \mathbb{N}^d$, the following estimate holds:*

$$\sum_{r=1}^{|\beta|} \sum_{p(\beta,r)} \frac{r!}{k_1! \cdot ... \cdot k_{|\beta|}!} \leq 2^{|\beta|(d+1)}$$

Proof. Let $f(\lambda) = \lambda^{|\beta|}$ and $g(x) = (-x_1)^{-1} \cdot ... \cdot (-x_d)^{-1}$. We apply the Faà di Bruno formula to the composition $g(f(x))$ at the point $x = (-1, ..., -1)$:

$$\begin{aligned} \prod_{\substack{j=1 \\ \beta_j \neq 0}}^{d} (|\beta|(|\beta|+1) \cdot ... \cdot (|\beta|+\beta_j-1)) &= \partial^\beta (g \circ f)(-1) \\ &= \sum_{r=1}^{|\beta|} \frac{|\beta|!}{(|\beta|-r)!} \sum_{p(\beta,r)} \beta! \prod_{j=1}^{|\beta|} \frac{1}{k_j!}. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} \prod_{\substack{j=1 \\ \beta_j \neq 0}}^{d} \frac{|\beta|(|\beta|+1) \cdot ... \cdot (|\beta|+\beta_j-1)}{\beta_j!} &= \sum_{r=1}^{|\beta|} \binom{|\beta|}{r} \sum_{p(\beta,r)} \frac{r!}{k_1! \cdot ... \cdot k_{|\beta|}!} \\ &\geq \sum_{r=1}^{|\beta|} \sum_{p(\beta,r)} \frac{r!}{k_1! \cdot ... \cdot k_{|\beta|}!}. \end{aligned}$$

From this, the desired inequality follows, since

$$\frac{|\beta|(|\beta|+1) \cdot ... \cdot (|\beta|+\beta_j-1)}{\beta_j!} = \frac{(|\beta|+\beta_j)!}{\beta_j!(|\beta|-1)!(|\beta|+\beta_j)} \leq \binom{|\beta|+\beta_j}{\beta_j} \leq 2^{|\beta|+\beta_j}.$$

□

If we apply the Faà di Bruno formula to the composition of a_0 and $w \mapsto \langle w \rangle$ and use the well known estimate $|\partial^\alpha \langle w \rangle| \leq C' 2^{|\alpha|+1} |\alpha|! \langle w \rangle^{1-|\alpha|}$, $\alpha \in \mathbb{N}^{2d}$, $w \in \mathbb{R}^{2d}$, we have

$$\begin{aligned} |D^\alpha a(w)| &\leq \sum_{r=1}^{|\alpha|} |(D^r a_0)(\langle w \rangle)| \sum_{p(\alpha,r)} \alpha! \prod_{j=1}^{|\alpha|} \frac{\left| D^{\alpha^{(j)}} \langle w \rangle \right|^{k_j}}{k_j! (\alpha^{(j)}!)^{k_j}} \\ &\leq \sum_{r=1}^{|\alpha|} \frac{h^{r/(l+l')} a(w)}{\langle w \rangle^{r(l+l'-1)/(l+l')}} \sum_{p(\alpha,r)} \alpha! \prod_{j=1}^{|\alpha|} \frac{C'^{k_j} 2^{|\alpha^{(j)}|k_j + k_j} \left(|\alpha^{(j)}|! \right)^{k_j} \langle w \rangle^{k_j - |\alpha^{(j)}|k_j}}{k_j! (\alpha^{(j)}!)^{k_j}} \\ &\leq \frac{2^{|\alpha|} C_0^{|\alpha|} a(w)}{\langle w \rangle^{|\alpha|(l+l'-1)/(l+l')}} \sum_{r=1}^{|\alpha|} \sum_{p(\alpha,r)} \alpha! \prod_{j=1}^{|\alpha|} \frac{\left(|\alpha^{(j)}|! \right)^{k_j}}{(\alpha^{(j)}!)^{k_j}} \prod_{j=1}^{|\alpha|} \frac{1}{k_j!}, \end{aligned}$$

where we denote by C_0 the quantity $2C'h^{1/(l+l')}$. One can easily prove that for $\beta^{(1)},...,\beta^{(n)} \in \mathbb{N}^{2d}$,

$$\frac{\left(\beta^{(1)}+...+\beta^{(n)}\right)!}{\beta^{(1)}!\cdot...\cdot\beta^{(n)}!} \leq \frac{\left|\beta^{(1)}+...+\beta^{(n)}\right|!}{\left|\beta^{(1)}\right|!\cdot...\cdot\left|\beta^{(n)}\right|!}.$$

In fact one easily verifies this inequality for $n=2$ and the general case follows by induction. We obtain

$$\alpha!\prod_{j=1}^{|\alpha|}\frac{\left(\left|\alpha^{(j)}\right|!\right)^{k_j}}{(\alpha^{(j)}!)^{k_j}} \leq |\alpha|!. \tag{2.2}$$

Using this estimate, together with Lemma 2.2, we conclude that

$$|D^\alpha a(w)| \leq \frac{\left(2^{2d+2}C_0\right)^{|\alpha|}|\alpha|!a(w)}{\langle w\rangle^{|\alpha|(l+l'-1)/(l+l')}}.$$

Hence, if we take $l'>0$ large enough such that $(l+l'-1)/(l+l') \geq \rho$, we obtain $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ and the growth condition $ii)$ of Definition 1.1 is satisfied for a. By the definition of a the lower bound $i)$ of Definition 1.1 trivially holds for $\tilde{M}_p = p!^{l+l'}$ and some $m>0$ in the (M_p) case (resp. for $\tilde{M}_p = p!^{l+l'/2}$ and $k_p = p^{l'/2}$ in the $\{M_p\}$ case). Hence a is elliptic.

Symbols given in the Introduction, correspond to $m = l+l'$ with suitable ρ. Actually, applying the Faà di Bruno formula to the composition of $\lambda \mapsto \lambda^{1/m}$ and $w \mapsto \langle w\rangle$, by the same technique as above using (2.2), Lemma 2.2 and the estimate $\left|\left(\lambda^{1/m}\right)^{(r)}\right| \leq r!\lambda^{-r+1/m}$, $\lambda>0$, one can prove $\left|D^\alpha\left(\langle w\rangle^{1/m}\right)\right| \leq C^{|\alpha|}|\alpha|!\langle w\rangle^{-|\alpha|+1/m}$. Applying again the Faà di Bruno formula to the composition of $\lambda \mapsto e^\lambda$ and $w \mapsto \langle w\rangle^{1/m}$ we obtain the estimate

$$\left|D^\alpha\left(e^{\langle w\rangle^{1/m}}\right)\right| \leq C^{|\alpha|}|\alpha|!e^{\langle w\rangle^{1/m}}\langle w\rangle^{-|\alpha|(m-1)/m}$$

which implies that such symbols can be considered in Theorem 1.2.

Example 2. An interesting nontrivial example of nonlinear term satisfying the assumptions of Theorem 1.2 can be given in the following way. Define p_β by

$$p_\beta(x) = \sum_{\alpha\in\mathbb{N}^d} c_{\alpha,\beta}x^{\alpha+\beta}$$

where $c_{\alpha,\beta}$ satisfy the following condition: for every $h>0$ there exists $C>0$ such that $|c_{\alpha,\beta}| \leq Ch^{|\alpha|+|\beta|}/\tilde{M}_{\alpha+\beta}$ (resp. $|c_{\alpha,\beta}| \leq Ch^{|\alpha|+|\beta|}/(\tilde{M}_{\alpha+\beta}\prod_{j=1}^{|\alpha|+|\beta|}k_j)$). By [14, Proposition 4.5.] and by simple calculation we obtain that $p_\beta(x) = R_\beta^{-1}P_\beta(x)$ satisfy the condition of Theorem 1.2, where $R_\beta = \prod_{j=1}^{|\beta|}r_j$ for some $(r_p) \in \mathfrak{R}$ (for example $r_p = p^\epsilon$, $\epsilon>0$) and P_β are entire functions which satisfy the estimate: for every $h>0$ there exists $C>0$ such that $|P_\beta(z)| \leq Ce^{\tilde{M}(h|z|)}$, $z\in\mathbb{C}^d$ (resp. $|P_\beta(z)| \leq Ce^{\tilde{N}_{k_p}(h|z|)}$, $z\in\mathbb{C}^d$), i.e. P_β are ultrapolynomials of class $\{\tilde{M}_p\}$ (resp. $\{\tilde{N}_p\}$, with

$\tilde{N}_p = \tilde{M}_p \prod_{j=1}^{p} k_j$). Interesting examples of this form are $F[u] = P(x)\cos u$ and $F[u] = P(x)e^{u^k}$, $k \in \mathbb{Z}_+$, where $P(x)$ is an ultrapolynomial with growth condition as stated above (to verify that these are indeed examples of the stated form one only needs to write $\cos u$ and e^{u^k} in power series in u). These examples fit well in our analysis since we presumed in Theorem 1.2 that $u \in H^s(\mathbb{R}^d)$ for some $s > d/2$, which implies that u is bounded over $\mathbb{R}^d$.

3 Pseudodifferential operators on $\mathcal{S}^*(\mathbb{R}^d), \mathcal{S}^{*\prime}(\mathbb{R}^d)$

In this section we recall some results contained in [23] and concerning the calculus for pseudodifferential operators with symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$. For the purposes of this paper we need to modify slightly some statements with respect to [23]. The proof of these new assertions is completely analogous to the original ones and do not deserve to be repeated here.

Now we recall the notion of asymptotic expansion for symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$, cf. [23, Definition 2].

Definition 3.1. *Let M_p and A_p be as in the definition of $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ and let $m_0 = 0, m_p = M_p/M_{p-1}, p \in \mathbb{Z}_+$. We denote by $FS^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ the space of all formal sums $\sum_{j\in\mathbb{N}} a_j$ such that for some $B > 0$, $a_j \in \mathcal{C}^\infty(int\, Q^c_{Bm_j})$ and satisfy the following condition: there exists $m > 0$ such that for every $h > 0$ (resp. there exists $h > 0$ such that for every $m > 0$) we have*

$$\sup_{j\in\mathbb{N}} \sup_{\alpha,\beta\in\mathbb{N}^d} \sup_{(x,\xi)\in Q^c_{Bm_j}} \frac{|D^\alpha_\xi D^\beta_x a_j(x,\xi)|\langle(x,\xi)\rangle^{\rho(|\alpha+\beta|+2j)} e^{-M(m|x|)-M(m|\xi|)}}{h^{|\alpha+\beta|+2j} A_\alpha A_\beta A_j^2} < \infty.$$

Notice that any symbol $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ can be regarded as an element $\sum\limits_{j\in\mathbb{N}} a_j$ of $FS^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ with $a_0 = a, a_j = 0$ for $j \geq 1$.

Definition 3.2. *A symbol $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ is equivalent to $\sum_{j\in\mathbb{N}} a_j \in FS^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ (we write $a \sim \sum_{j\in\mathbb{N}} a_j$ in this case) if there exist $m, B > 0$ such that for every $h > 0$ (resp. there exist $h, B > 0$ such that for every $m > 0$) the following condition holds:*

$$\sup_{N\in\mathbb{Z}_+} \sup_{\alpha,\beta\in\mathbb{N}^d} \sup_{(x,\xi)\in Q^c_{Bm_N}} \frac{\left|D^\alpha_\xi D^\beta_x \big(a(x,\xi) - \sum\limits_{j<N} a_j(x,\xi)\big)\right| e^{-M(m|x|)-M(m|\xi|)}}{h^{|\alpha+\beta|+2N} A_\alpha A_\beta A_N^2 \langle(x,\xi)\rangle^{-\rho(|\alpha+\beta|+2N)}} < \infty.$$

An operator $a(x,D)$ with symbol $a \sim 0$ is $*$-regularizing, namely it extends to a linear and continuous map from $\mathcal{S}^{*\prime}(\mathbb{R}^d)$ to $\mathcal{S}^*(\mathbb{R}^d)$, see [23, Theorem 3]. Moreover, for every sum $\sum_{j\in\mathbb{N}} a_j \in FS^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ one can find a symbol $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ such that $a \sim \sum_{j\in\mathbb{N}} a_j$, cf. [23, Theorem 4]. Actually by the same argument one can prove the following more precise assertion.

Proposition 3.3. *Let g be a positive continuous function such that $g(w)$ and $1/g(w)$ have ultrapolynomial growth of class $*$ and let U be a subset of $FS^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ for which*

there exists $B > 0$ such that for every $h > 0$ there exists $C > 0$ (resp. there exist $B, h, C > 0$) such that

$$\sup_{j\in\mathbb{N}}\sup_{\alpha,\beta}\sup_{(x,\xi)\in Q^c_{Bm_j}}\frac{\left|D_\xi^\alpha D_x^\beta a_j(x,\xi)\right|\langle(x,\xi)\rangle^{\rho|\alpha|+\rho|\beta|+2j\rho}}{h^{|\alpha|+|\beta|+2j}A_\alpha A_\beta A_j^2 g(x,\xi)}\leq C$$

for all $\sum_{j\in\mathbb{N}} a_j \in U$. Then for every $\tilde{h} > 0$ there exists $C > 0$ (resp. there exist $\tilde{h}, C > 0$) such that the following condition holds: for every sum $\sum a_j \in U$ there exists a symbol $a \sim \sum_{j\in\mathbb{N}} a_j$ satisfying the following estimate:

$$\left|D_\xi^\alpha D_x^\beta a(x,\xi)\right| \leq C\frac{\tilde{h}^{|\alpha|+|\beta|}A_\alpha A_\beta g(x,\xi)}{\langle(x,\xi)\rangle^{\rho(|\alpha|+|\beta|)}}.$$

In [23, Theorem 7] it was proved that the composition of two operators $b(x,D)$ and $a(x,D)$ with symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ is the sum of an operator $f_{a,b}(x,D)$ with symbol $f_{a,b} \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$, with $f_{a,b} \sim \sum_j f_{a,b,j}$, where

$$f_{a,b,j}(x,\xi) = \sum_{|\alpha|=j}\frac{1}{\alpha!}\partial_\xi^\alpha b(x,\xi)D_x^\alpha a(x,\xi), \tag{3.1}$$

and of a *-regularizing operator $T_{a,b}$. In fact, $f_{a,b} = \sum_j(1-\chi_j)f_{a,b,j}$ where χ_j is defined in the following way (cf. the proof of [23, Theorem 4]). Take $\varphi, \psi \in \mathcal{D}^{(A_p)}\left(\mathbb{R}^d\right)$, in the (M_p) case, resp. $\varphi, \psi \in \mathcal{D}^{\{A_p\}}\left(\mathbb{R}^d\right)$ in the $\{M_p\}$ case, such that $0 \leq \varphi, \psi \leq 1$, $\varphi(x) = 1$ when $\langle x\rangle \leq 2$, $\psi(\xi) = 1$ when $\langle\xi\rangle \leq 2$ and $\varphi(x) = 0$ when $\langle x\rangle \geq 3$, $\psi(\xi) = 0$ when $\langle\xi\rangle \geq 3$. Then define $\chi(x,\xi) = \varphi(x)\psi(\xi)$, $\chi_n(x,\xi) = \chi\left(\frac{x}{Rm_n}, \frac{\xi}{Rm_n}\right)$ for $n \in \mathbb{Z}_+$ and $\chi_0(x,\xi) = 0$, where $m_n = M_n/M_{n-1}$ and $R > 0$ is large enough.

Proposition 3.4. *Let U_1 and U_2 be bounded subsets of $\Gamma^{(M_p),\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}, m'\right)$ (resp. $\Gamma^{\{M_p\},\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}, h'\right)$), for some $m' > 0$ (resp. for some $h' > 0$). Then for every $a \in U_1$ and $b \in U_2$ we have $b(x,D)a(x,D) = f_{a,b}(x,D) + T_{a,b}$ where $f_{a,b} = \sum_j(1-\chi_j)f_{a,b,j}$ and χ_j are the cut-off functions defined above which can be chosen uniformly for $a \in U_1, b \in U_2$, and with $f_{a,b,j}$ given by (3.1). Moreover, the family $T_{a,b}$ of *-regularizing operators is an equicontinuous subset of $\mathcal{L}_b\left(\mathcal{S}'^*\left(\mathbb{R}^d\right), \mathcal{S}^*\left(\mathbb{R}^d\right)\right)$.*

From the results above we notice that in general the composition of two operators with symbols in $\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ is still an operator of infinite order. In the sequel we will be interested to the case when the composition is a finite order operator with bounded symbol, hence the related operator is bounded on Sobolev spaces. With this purpose we give the following definition.

Definition 3.5. *Let $V, W \subseteq \Gamma^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ and let $f(w)$ be a positive continuous function on $\mathbb{R}^{2d}$ such that $f(w)$ and $1/f(w)$ are of ultrapolynomial growth of class * (see [16]). The sets V and W are said to be $(f,*)$-conjugate if for every $h > 0$ there exists $C > 0$ (resp. there exist $h, C > 0$) such that*

$$|D^\alpha a(w)| \leq C\frac{h^{|\alpha|}A_\alpha}{\langle w\rangle^{\rho|\alpha|}f(w)} \text{ and } |D^\alpha b(w)| \leq C\frac{h^{|\alpha|}A_\alpha f(w)}{\langle w\rangle^{\rho|\alpha|}} \text{ for all } a \in V,\ b \in W.$$

Obviously if V and W are $(f,*)$-conjugate then they are bounded subsets of $\Gamma^{(M_p),\infty}_{A_p,\rho}\left(\mathbb{R}^{2d};m'\right)$ for some $m'>0$ (resp. $\Gamma^{\{M_p\},\infty}_{A_p,\rho}\left(\mathbb{R}^{2d};h'\right)$, for some $h'>0$).

Proposition 3.6. *Let V and W be $(f,*)$-conjugate. Then, there exists $C>0$ such that*

$$\|b(x,D)a(x,D)\|_{\mathcal{L}_b(H^s)}\leq C,\ \textit{for all } a\in V,\, b\in W.$$

Proof. Let $f_{a,b}$ be the symbol of the operator $b(x,D)a(x,D)$ defined as above. Then $b(x,D)a(x,D)=f_{a,b}(x,D)+T_{a,b}$, where $T_{a,b}$ form an equicontinuous subset of $\mathcal{L}_b\left(\mathcal{S}'^*\left(\mathbb{R}^d\right),\mathcal{S}^*\left(\mathbb{R}^d\right)\right)$. Then $f_{a,b}\sim\sum_j f_{a,b,j}$, where

$$f_{a,b,j}(w)=\sum_{|\nu|=j}\frac{1}{\nu!}\partial_\xi^\nu b(w)D_x^\nu a(w).$$

Observe that

$$\begin{aligned}|D_w^\alpha f_{a,b,j}(w)| &\leq \sum_{\beta\leq\alpha}\sum_{|\nu|=j}\frac{1}{\nu!}\left|D_w^{\alpha-\beta}D_\xi^\nu b(w)\right|\left|D_w^\beta D_x^\nu a(w)\right|\\ &\leq C_1\sum_{\beta\leq\alpha}\sum_{|\nu|=j}\frac{1}{\nu!}\frac{h^{|\alpha|+2|\nu|}A_{|\alpha|+2|\nu|}}{\langle w\rangle^{\rho(|\alpha|+2|\nu|)}}\leq C_2\frac{(2hH)^{|\alpha|+2j}A_\alpha A_{2j}}{\langle w\rangle^{\rho(|\alpha|+2j)}}.\end{aligned}$$

Now, since $f_{a,b}=\sum_j(1-\chi_j)f_{a,b,j}$ with χ_j defined as above, one easily obtains that for every $h>0$ there exists $C>0$, resp. there exist $h,C>0$ such that $|D_w^\alpha f_{a,b}(w)|\leq Ch^{|\alpha|}A_\alpha\langle w\rangle^{-\rho|\alpha|}$. From this it follows that $f_{a,b}(x,D)$, $a\in V$, $b\in W$, form a bounded subset of $\mathcal{L}_b(H^s)$ (cf. [18, Theorem 1.7.14] and [18, Theorem 2.1.11] and its proof), the claim follows. □

The next result has been proved in [8] for more general hypoelliptic operators. It is immediate to verify that it holds in particular for symbols satisfying the ellipticity conditions in Definition 1.1.

Theorem 3.7. *Let $a\in\Gamma^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic). Then there exists a *-regularizing operator T and a symbol $b\in\Gamma^{*,\infty}_{A_p,\rho}\left(\mathbb{R}^{2d}\right)$ such that $b(x,D)a(x,D)=\mathrm{Id}+T$. Moreover, the symbol b satisfies the following condition: there exists $B'>0$ such that for every $h>0$ there exist $C>0$ (resp. there exist $h,C>0$) such that*

$$\left|D_\xi^\alpha D_x^\beta b(x,\xi)\right|\leq C\frac{h^{|\alpha|+|\beta|}A_{\alpha+\beta}}{|a(x,\xi)|\langle(x,\xi)\rangle^{\rho(|\alpha|+|\beta|)}},\qquad(x,\xi)\in Q^c_{B'}.\tag{3.2}$$

As we mentioned in the introduction, given $(r_p)\in\mathfrak{R}$ and multi-index γ, R_γ stands for $R_{|\gamma|}=\prod_{j=1}^{|\gamma|}r_j$.

Lemma 3.8. *Let $a\in\Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) and let b be the symbol of the parametrix of $a(x,D)$. Then the sets $\{b\}$ and $\left\{\dfrac{h^{|\alpha|}}{M_\alpha}D_w^\alpha a\,\middle|\,\alpha\in\mathbb{N}^{2d}\right\}$*

are $(|a(w)|,*)$-conjugate for every $h>0$ (resp. for some $h>0$). Hence, for every $h>0$ there exists $C>0$ (resp. there exist $h,C>0$) such that the estimate $\|b(x,D)\circ(D_w^\alpha a)(x,D)\|_{\mathcal{L}_b(H^s)}\leq Ch^{|\alpha|}M_\alpha$ holds. Moreover, in the (M_p) case, there exist $(r_p)\in\mathfrak{R}$ with $r_1=1$ and $C>0$ such that $\|b(x,D)\circ(D_w^\alpha a)(x,D)\|_{\mathcal{L}_b(H^s)}\leq CM_\alpha/R_\alpha$.

Proof. We have

$$\begin{aligned}\frac{h^{|\alpha|}}{M_\alpha}\left|D_w^{\alpha+\beta}a(w)\right| &\leq C_1\frac{h^{|\alpha|}h_1^{|\alpha|+|\beta|}A_{\alpha+\beta}|a(w)|}{M_\alpha\langle w\rangle^{\rho(|\alpha|+|\beta|)}}\\ &\leq C_2\frac{(LHhh_1)^{|\alpha|}(Hh_1)^{|\beta|}A_\beta|a(w)|}{\langle w\rangle^{\rho|\beta|}}.\end{aligned}$$

The $\{M_p\}$ case trivially follows from this by choosing h small enough, since h_1 is fixed. In the (M_p) case, for each fixed h we can take h_1 arbitrary small. One easily sees that this implies that the sets under consideration are $(|a(w)|,(M_p))$-conjugate. The inequality $\|b(x,D)\circ(D_w^\alpha a)(x,D)\|_{\mathcal{L}_b(H^s)}\leq Ch^{|\alpha|}M_\alpha$ follows by Proposition 3.6. It remains to prove the last part of the lemma. Since, in the (M_p) case, we already proved that $\sup_{\alpha\in\mathbb{N}^{2d}}\dfrac{\|b(x,D)\circ(D_w^\alpha a)(x,D)\|_{\mathcal{L}_b(H^s)}}{h^{|\alpha|}M_\alpha}<\infty$ for every $h>0$, by Lemma 3.4 of [17] we can conclude that there exists $(\tilde r_p)\in\mathfrak{R}$ and $C>0$ such that $\|b(x,D)\circ(D_w^\alpha a)(x,D)\|_{\mathcal{L}_b(H^s)}\leq CM_\alpha/\prod_{j=1}^{|\alpha|}\tilde r_j$. If we take $r_p=\max\{\tilde r_p,1\}$, then $(r_p)\in\mathfrak{R}$, $r_1=1$ and the desired estimate holds for this (r_p), possibly with a larger constant C. □

Lemma 3.9. *There exists $l\geq1$ such that the sets $\left\{\dfrac{h^{|\beta|}}{l^{|\gamma|}\tilde M_\gamma}D_x^\alpha p_\beta\xi^\gamma\,\middle|\,\alpha,\beta,\gamma\in\mathbb{N}^d\right\}$ and $\{b\}$ are $\left(e^{\tilde M(m|x|)}e^{\tilde M(m|\xi|)},(M_p)\right)$-conjugate for any $h>0$, (resp. the sets $\left\{\dfrac{h^{|\beta|}}{l^{|\gamma|}K_\gamma\tilde M_\gamma}D_x^\alpha p_\beta\xi^\gamma\,\middle|\,\alpha,\beta,\gamma\in\mathbb{N}^d\right\}$ and $\{b\}$ are $\left(e^{\tilde N_{k_p}(|x|)}e^{\tilde N_{k_p}(|\xi|)},\{M_p\}\right)$-conjugate for any $h>0$). In particular for each $\tilde h>0$ there exists $C>0$ such that $\|(B\circ D^\alpha p_\beta(x)\partial^\gamma)(x,D)\|_{\mathcal{L}(H^s)}\leq C\tilde h^{|\beta|}l^{|\gamma|}\tilde M_\gamma$ (resp. $\|(B\circ D^\alpha p_\beta(x)\partial^\gamma)(x,D)\|_{\mathcal{L}(H^s)}\leq C\tilde h^{|\beta|}l^{|\gamma|}K_\gamma\tilde M_\gamma$).*

Proof. We consider first the (M_p) case. Pick $l\geq1$ such that $H^2/l\leq m/12$. Let $h,h'>0$ be arbitrary but fixed. Pick $0<h_1<1$ such that $H\sqrt{h_1}\leq h'$, $L\tilde LHh_1\leq1$, $hh_1\leq1$ and $H^2\sqrt{h_1}\leq m/6$. Then

$$\begin{aligned}&\frac{h^{|\beta|}}{l^{|\gamma|}\tilde M_\gamma}\left|D_\xi^\mu D_x^\nu\left(D_x^\alpha p_\beta(x)\xi^\gamma\right)\right|\\ &\leq \frac{h^{|\beta|}\gamma!}{l^{|\gamma|}\tilde M_\gamma(\gamma-\mu)!}|\xi|^{|\gamma|-|\mu|}\left|D_x^{\alpha+\nu}p_\beta(x)\right|\\ &\leq C_1\frac{2^{|\gamma|}h^{|\beta|}h_1^{|\alpha|+|\beta|+|\nu|}A_{\alpha+\nu}\mu!\langle(x,\xi)\rangle^{|\mu|+|\nu|}}{l^{|\gamma|}\tilde M_\gamma\tilde M_{\alpha+\nu}\langle(x,\xi)\rangle^{\rho(|\mu|+|\nu|)}}|\xi|^{|\gamma|-|\mu|}e^{\tilde M(h_1|x|)}\end{aligned}$$

$$\leq C_2\frac{(2H/l)^{|\gamma|}(H^2h_1)^{|\nu|}(Hh_1)^{|\alpha|}(hh_1)^{|\beta|}A_\alpha A_\nu \mu!\langle(x,\xi)\rangle^{|\gamma|+|\nu|}}{\tilde{M}_{\gamma+\nu}\tilde{M}_\alpha\langle(x,\xi)\rangle^{\rho(|\mu|+|\nu|)}}e^{\tilde{M}(h_1|x|)}$$
$$\leq C_3\frac{(H\sqrt{h_1})^{|\nu|}(L\tilde{L}Hh_1)^{|\alpha|}(hh_1)^{|\beta|}h'^{|\mu|}A_{\nu+\mu}}{\langle(x,\xi)\rangle^{\rho(|\mu|+|\nu|)}}e^{\tilde{M}(h_1|x|)}e^{\tilde{M}((H\sqrt{h_1}+2H/l)\langle(x,\xi)\rangle)}.$$

Since $\tilde{M}_p$ satisfies $(M.2)$, by Proposition 3.6 of [14], we have

$$\begin{aligned}e^{\tilde{M}(h_1|x|)}e^{\tilde{M}((H\sqrt{h_1}+2H/l)\langle(x,\xi)\rangle)} &\leq C_4e^{\tilde{M}(h_1|x|)}e^{\tilde{M}(3(H\sqrt{h_1}+2H/l)|x|)}e^{\tilde{M}(3(H\sqrt{h_1}+2H/l)|\xi|)}\\ &\leq C_5e^{\tilde{M}(3(H\sqrt{h_1}+2H/l)|\xi|)}e^{\tilde{M}(3(H^2\sqrt{h_1}+2H^2/l)|x|)}.\end{aligned}$$

If we use this in the above estimate, by the way we defined h_1, we have

$$\frac{h^{|\beta|}}{l^{|\gamma|}\tilde{M}_\gamma}\left|D_\xi^\mu D_x^\nu\left(D_x^\alpha p_\beta(x)\xi^\gamma\right)\right| \leq C\frac{h'^{|\mu|+|\nu|}A_{\mu+\nu}}{\langle(x,\xi)\rangle^{\rho(|\mu|+|\nu|)}}e^{\tilde{M}(m|\xi|)}e^{\tilde{M}(m|x|)},$$

which proofs the (M_p) case. In the $\{M_p\}$ case one can use the same technique as above (observe that the sequence $K_p\tilde{M}_p$ satisfies $(M.2)$). The last part follows by Proposition 3.6. □

Lemma 3.10. *Let $h>0$ and for each $\beta\in\mathbb{N}^d$, let $p_\beta(x)$ be a smooth function satisfying (1.3) in the (M_p) case (resp. satisfying (1.4) in the $\{M_p\}$ case) and let $j_\beta\in\{1,...,d\}$. Then the following properties hold:*

a) The sets $\{b\}$ and $\left\{h^{|\beta|}x_{j_\beta}p_\beta(x)\big|\,\beta\in\mathbb{N}^d\right\}$ are $\left(e^{\tilde{M}(m|x|)}e^{\tilde{M}(m|\xi|)},(M_p)\right)$-conjugate (resp. $\left(e^{\tilde{N}_{k_p}(|x|)}e^{\tilde{N}_{k_p}(|\xi|)},\{M_p\}\right)$-conjugate). In particular, for every $\tilde{h}>0$ there exists $C>0$, such that $\left\|(B\circ x_{j_\beta}p_\beta(x))(x,D)\right\|_{\mathcal{L}_b(H^s)}\leq C\tilde{h}^{|\beta|}$. Moreover, there exist $(r_p)\in\mathfrak{R}$ with $r_1=1$ and $C>0$ such that $\left\|(B\circ x_{j_\beta}p_\beta(x))(x,D)\right\|_{\mathcal{L}_b(H^s)}\leq C/R_\beta$.

b) The sets $\{b\}$ and $\left\{h^{|\beta|}\xi_jp_\beta(x)\big|\,\beta\in\mathbb{N}^d\right\}$ are $\left(e^{\tilde{M}(m|x|)}e^{\tilde{M}(m|\xi|)},(M_p)\right)$-conjugate (resp. $\left(e^{\tilde{N}_{k_p}(|x|)}e^{\tilde{N}_{k_p}(|\xi|)},\{M_p\}\right)$-conjugate). In particular, for every $\tilde{h}>0$ there exists $C>0$ such that $\|(B\circ p_\beta(x)\partial_j)(x,D)\|_{\mathcal{L}_b(H^s)}\leq C\tilde{h}^{|\beta|}$. Moreover, there exist $(r_p)\in\mathfrak{R}$ with $r_1=1$ and $C>0$ such that $\|(B\circ p_\beta(x)\partial_j)(x,D)\|_{\mathcal{L}_b(H^s)}\leq C/R_\beta$.

Proof. We prove $a)$, the proof of $b)$ being completely analogous. In the (M_p) case, let $h,h'>0$ be arbitrary but fixed. We have

$$\begin{aligned}h^{|\beta|}\left|D^\alpha\left(x^{e_{j_\beta}}p_\beta(x)\right)\right| &\leq h^{|\beta|}\left|x_{j_\beta}\right|\left|D^\alpha p_\beta(x)\right|+|\alpha|h^{|\beta|}\left|D^{\alpha-e_{j_\beta}}p_\beta(x)\right|\\ &= S_1(x)+S_2(x).\end{aligned}$$

Take $h_2<1$ such that $3h_2H\leq m/2$ and take $h_1<1$ such that $2Hh_1/h_2\leq h'$, $Hh_1\leq m/2$ and $h_1\leq 1/h$. To estimate $S_1(x)$ we have

$$\begin{aligned}S_1(x) &\leq C_1h^{|\beta|}|x|\frac{(Hh_1)^{|\alpha|}h_1^{|\beta|}A_\alpha e^{\tilde{M}(h_1|x|)}\langle(x,\xi)\rangle^{|\alpha|}}{\tilde{M}_{|\alpha|+1}\langle(x,\xi)\rangle^{\rho|\alpha|}}\\ &\leq C_1\frac{(Hh_1)^{|\alpha|}A_\alpha e^{\tilde{M}(h_1|x|)}e^{\tilde{M}(h_2\langle(x,\xi)\rangle)}}{h_2^{|\alpha|+1}\langle(x,\xi)\rangle^{\rho|\alpha|}}\end{aligned}$$

$$\leq \quad C_2 \frac{h'^{|\alpha|} A_\alpha e^{\tilde{M}((h_1+3h_2)H|x|)} e^{\tilde{M}(3h_2|\xi|)}}{\langle (x,\xi)\rangle^{\rho|\alpha|}} \leq C_2 \frac{h'^{|\alpha|} A_\alpha e^{\tilde{M}(m|\xi|)} e^{\tilde{M}(m|x|)}}{\langle (x,\xi)\rangle^{\rho|\alpha|}}.$$

Similar estimates can be obtained for $S_2(x)$ in the same way and the $\{M_p\}$ case can be treated similarly. The estimate $\|(B \circ x_j p_\beta(x))(x,D)\|_{\mathcal{L}_b(H^s)} \leq C\tilde{h}^{|\beta|}$ follows from Proposition 3.6. The last part can be proved similarly as in the proof of Lemma 3.8, by using Lemma 3.4 of [17]. □

4 The proof of Theorem 1.2

The proof of Theorem 1.2 needs some preparation. First of all it is useful to characterize the space $\mathcal{S}^*(\mathbb{R}^d)$ in terms of suitable scales of Sobolev norms.

Namely, let

$$\|\varphi\|_{s,h} = \sum_{\alpha\in\mathbb{N}^d} \frac{h^{|\alpha|}}{M_\alpha} \|x^\alpha \varphi(x)\|_{H^s} \quad \text{and} \quad \|\varphi\|_{\{s,h\}} = \sum_{\alpha\in\mathbb{N}^d} \frac{h^{|\alpha|}}{M_\alpha} \|D^\alpha \varphi(x)\|_{H^s}.$$

Moreover, for $h > 0$ and $(r_p) \in \mathfrak{R}$, set

$$H_N^{s,h}[\varphi] = \sum_{|\alpha|\leq N} \frac{h^{|\alpha|}}{M_\alpha} \|x^\alpha \varphi(x)\|_{H^s} \quad , \quad H_N^{s,h,(r_p)}[\varphi] = \sum_{|\alpha|\leq N} \frac{h^{|\alpha|} R_\alpha}{M_\alpha} \|x^\alpha \varphi(x)\|_{H^s},$$

$$E_N^{s,h}[\varphi] = \sum_{|\alpha|\leq N} \frac{h^{|\alpha|}}{M_\alpha} \|D^\alpha \varphi(x)\|_{H^s} \quad , \quad E_N^{s,h,(r_p)}[\varphi] = \sum_{|\alpha|\leq N} \frac{h^{|\alpha|} R_\alpha}{M_\alpha} \|D^\alpha \varphi(x)\|_{H^s}.$$

Let $\varphi \in \mathcal{S}\left(\mathbb{R}^d\right)$. We recall the well known result (see for example [9] or [18]).

Lemma 4.1. *The following conditions are equivalent:*

i) $\varphi \in \mathcal{S}^*\left(\mathbb{R}^d\right)$;

ii) there exists $s > d/2$ *such that for every* $h > 0$ *(resp. there exists* $h > 0$*) such that* $\|\varphi\|_{s,h} < \infty$ *and* $\|\varphi\|_{\{s,h\}} < \infty$.

By Lemma 4.1 we can prove that a function $u \in \mathcal{S}^*(\mathbb{R}^d)$ by proving the decay and the regularity properties separately. This allows to simplify considerably the proofs, see also [5, 6].

Next we state a preliminary technical result which will be used in the subsequent proofs.

Lemma 4.2. *Let* M_p *be a sequence which satisfies* $(M.3)'$, $(M.4)$ *and* $M_0 = M_1 = 1$. *Let* $(k'_p), (k''_p) \in \mathfrak{R}$, $k'_1 = k''_1 = 1$. *There exists* $(r'_p) \in \mathfrak{R}$ *such that* $r'_1 = 1$, $(r'_p) \leq (k'_p)$, $(r'_p) \leq (k''_p)$ *and the sequence* $N_p = M_p / \prod_{j=1}^p r'_p$, *for* $p \in \mathbb{Z}_+$ *and* $N_0 = 1$ *satisfies* $(M.3)'$ *and* $(M.4)$.

Proof. Let $a_p > 0$, $p \in \mathbb{Z}_+$, are such that $\sum_{p=1}^\infty a_p$ is convergent. Then one easily verifies that $\sum_{p=1}^\infty \frac{a_p}{s_p}$ is also convergent, where $s_p = \sqrt{\sum_{j=p}^\infty a_j}$, $p \in \mathbb{Z}_+$ (one easily

obtains that the partial sums of the series $\sum a_p/s_p$ are a Cauchy sequence). Put $\tilde{c} = \sqrt{\sum_{j=1}^{\infty} 1/m_j}$ and define $\tilde{r}'_p = \tilde{c}\left(\sum_{j=p}^{\infty} \frac{1}{m_j}\right)^{-1/2}$, $p \in \mathbb{Z}_+$. Then we obtain that $\tilde{r}'_1 = 1$, $(\tilde{r}'_p) \in \mathfrak{R}$ and $\sum \tilde{r}'_p/m_p$ converges. Let $r_p = \min\{k'_p, k''_p, \tilde{r}'_p\}$, for $p \in \mathbb{Z}_+$. Then, obviously, $r_1 = 1$, $(r_p) \in \mathfrak{R}$, $(r_p) \leq (k'_p)$, $(r_p) \leq (k''_p)$ and $(r_p) \leq (\tilde{r}'_p)$. Also $\sum r_p/m_p$ converges. Define the sequence (r'_p) by $r'_1 = 1$ and inductively

$$r'_{p+1} = \min\left\{r_{p+1}, \frac{pm_{p+1}}{(p+1)m_p} r'_p\right\},$$

for $p \in \mathbb{Z}_+$. We will prove that this (r'_p) satisfies the desired conditions. First, note that $r'_p \leq r_p$, for all $p \in \mathbb{Z}_+$. Since $r_{p+1} \geq r_p$ and $pm_{p+1} \geq (p+1)m_p$ (which is equivalent to $(M.4)$ for M_p) it follows that

$$r'_{p+1} = \min\left\{r_{p+1}, \frac{pm_{p+1}}{(p+1)m_p} r'_p\right\} \geq \min\{r_p, r'_p\} = r'_p,$$

for all $p \in \mathbb{Z}_+$. To prove that r'_p tends to infinity, assume the contrary. Since we already proved that r'_p is monotonically increasing, there exists $C > 1$ such that $r'_p \leq C$ for all $p \in \mathbb{Z}_+$. Since $(r_p) \in \mathfrak{R}$, there exists $p_0 \in \mathbb{Z}_+$ such that $r_p > C + 1$ for all $p \geq p_0$. But then, $r'_{p+1} = \frac{pm_{p+1}}{(p+1)m_p} r'_p$ for all $p \geq p_0$. Then, for $p \geq p_0$, we have

$$\begin{aligned} r'_{p+1} &= \frac{pm_{p+1}}{(p+1)m_p} r'_p = \frac{pm_{p+1}}{(p+1)m_p} \cdot \frac{(p-1)m_p}{pm_{p-1}} r'_{p-1} \\ &= \ldots = \frac{pm_{p+1}}{(p+1)m_p} \cdot \frac{(p-1)m_p}{pm_{p-1}} \cdot \ldots \cdot \frac{p_0 m_{p_0+1}}{(p_0+1)m_{p_0}} r'_{p_0} \\ &= \frac{p_0 m_{p+1}}{(p+1)m_{p_0}} r'_{p_0} \end{aligned}$$

which tends to infinity when $p \to \infty$ because of $(M.3)'$ for M_p. Hence $(r'_p) \in \mathfrak{R}$. The claim that N_p satisfies $(M.4)$ is equivalent to $r'_{p+1} \leq \frac{pm_{p+1}}{(p+1)m_p} r'_p$, which is true by the way we defined the sequence (r'_p). Moreover, if we put $n_p = N_p/N_{p-1}$, then $n_p = m_p/r'_p \geq m_p/r_p$ and we know that $\sum r_p/m_p$ converges. Hence N_p satisfies $(M.3)'$. □

After these preliminaries we can prove the following two results.

Theorem 4.3. *Let $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) and let $f \in \mathcal{S}^*(\mathbb{R}^d)$. Assume that $u \in H^s(\mathbb{R}^d)$, $s > d/2$, is a solution of $Au = f + F[u]$, where $F[u]$ is defined by (1.3) and (1.5) (resp. (1.4) and (1.5)). Then we have $\|u\|_{s,h} < \infty$ for every $h > 0$ (resp. for some $h > 0$).*

Theorem 4.4. *Let $a \in \Gamma^{*,\infty}_{A_p,\rho}(\mathbb{R}^{2d})$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) and let $f \in \mathcal{S}^*(\mathbb{R}^d)$. Assume that $u \in H^s(\mathbb{R}^d)$, $s > d/2$, is a solution of $Au = f + F[u]$, where $F[u]$ is defined by (1.3) and (1.5) (resp. (1.4) and (1.5)).*

i) If $F[u]$ is a finite sum, then we have $\|u\|_{\{s,h\}} < \infty$ for every $h > 0$ (resp. for some $h > 0$).

ii) If $F[u]$ is infinite sum, then $\sum_{\alpha} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|\partial^\alpha u\|_{H^s} < \infty$ for some $h > 0$ in the (M_p) case (resp. $\sum_{\alpha} \frac{h^{|\alpha|}\|\partial^\alpha u\|_{H^s}}{\tilde{M}_\alpha \prod_{j=1}^{|\alpha|} k_j} < \infty$ for some $h > 0$ in the $\{M_p\}$ case).

Notice that by Lemma 4.1, Theorem 1.2 follows directly from the combination of Theorems 4.3 and 4.4. Let us prove the two latter results.

Lemma 4.5. *Let $A = a(x, D)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) operator and let B be its parametrix. Then the following properties hold:*

i) In the (M_p) case, let $(r_p) \in \mathfrak{R}$ be the sequence in Lemma 3.8. Let $(r'_p) \in \mathfrak{R}$ be a sequence such that $(r'_p) \leq (r_p)$, $r'_1 = 1$ and the sequence M_p/R'_p satisfies $(M.3)'$ and $(M.4)$. Then for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for every $0 < h < h_0$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|} R'_\alpha}{M_\alpha} \|B[A, x^\alpha]u\|_{H^s} \leq \varepsilon H_{N-1}^{s,h,(r'_p)}[u].$$

ii) In the $\{M_p\}$ case, for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for all $0 < h < h_0(\varepsilon)$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{M_\alpha} \|B[A, x^\alpha]u\|_{H^s} \leq \varepsilon H_{N-1}^{s,h}[u].$$

Proof. First we prove the (M_p) case. (The existence of such sequence $(r'_p) \in \mathfrak{R}$ is given by Lemma 4.2.) For shorter notation, put $N_p = M_p/R'_p$, for $p \in \mathbb{Z}_+$ and $N_0 = 1$. Observe that

$$x^\alpha A u(x) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (-1)^{|\beta|} (D_\xi^\beta a)(x, D)(x^{\alpha-\beta} u(x)).$$

So, we obtain

$$B[A, x^\alpha]u = \sum_{\substack{\beta \leq \alpha \\ \beta \neq 0}} \binom{\alpha}{\beta} (-1)^{|\beta|+1} B(D_\xi^\beta a)(x, D)(x^{\alpha-\beta} u(x)).$$

By Lemma 3.8, there exists $C > 0$ such that $\left\| B(D_\xi^\beta a)(x, D) \right\|_{\mathcal{L}_b(H^s)} \leq C N_\beta$. Let $0 < \varepsilon < 1$ be fixed. Choose $0 < h_0 < 1/2$ such that $h_0 < \varepsilon \left(2C \sum_{|\beta|=1}^{\infty} 2^{-|\beta|+1} \right)^{-1}$. For $0 < h < h_0$ we obtain

$$
\begin{aligned}
&\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B[A, x^\alpha]u\|_{H^s} \\
&\leq \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \sum_{\substack{\beta\leq\alpha \\ \beta\neq 0}} \binom{\alpha}{\beta} \left\| B(D_\xi^\beta a)(x,D) x^{\alpha-\beta} u \right\|_{H^s} \\
&\leq C \sum_{|\alpha|=1}^{N} \sum_{\substack{\beta\leq\alpha \\ \beta\neq 0}} \frac{h^{|\alpha|} N_\beta}{N_\alpha} \binom{\alpha}{\beta} \left\| x^{\alpha-\beta} u \right\|_{H^s} \leq C \sum_{|\beta|=1}^{N} h^{|\beta|} \sum_{\substack{\alpha\geq\beta \\ |\alpha|\leq N}} \frac{h^{|\alpha|-|\beta|}}{N_{\alpha-\beta}} \left\| x^{\alpha-\beta} u \right\|_{H^s} \\
&\leq \varepsilon H_{N-1}^{s,h,(r'_p)}[u],
\end{aligned}
$$

where in the third inequality, we used $(M.4)$ for N_p and the fact $\binom{\alpha}{\beta} \leq \binom{|\alpha|}{|\beta|}$. This completes the proof in the (M_p) case. For the $\{M_p\}$ case, let $\varepsilon > 0$. By Lemma 3.8, there exist $h_1, C > 0$ such that $\left\| B(D_\xi^\beta a)(x,D) \right\|_{\mathcal{L}_b(H^s)} \leq C h_1^{|\beta|} M_\beta$. Choose $h_0 > 0$ such that $h_0 h_1 < 1/2$ and $h_0 h_1 \leq \varepsilon \left(2C \sum_{|\beta|=1}^{\infty} 2^{-|\beta|+1} \right)^{-1}$. Then, for $0 < h < h_0$, similarly as before, we obtain

$$
\begin{aligned}
\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{M_\alpha} \|B[A, x^\alpha]u\|_{H^s} &\leq C \sum_{|\beta|=1}^{N} (h h_1)^{|\beta|} \sum_{\substack{\alpha\geq\beta \\ |\alpha|\leq N}} \frac{h^{|\alpha|-|\beta|}}{M_{\alpha-\beta}} \left\| x^{\alpha-\beta} u \right\|_{H^s} \\
&\leq \varepsilon H_{N-1}^{s,h}[u],
\end{aligned}
$$

which completes the proof. □

Lemma 4.6. *Let $A = a(x,D)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) operator and let B be its parametrix. Let $F[u]$ be defined by (1.3), (1.5) in the (M_p) case (resp. by (1.4), (1.5) in the $\{M_p\}$ case). Then the following properties hold:*

i) In the (M_p) case, let $(r_p) \in \mathfrak{R}$ be the sequence in Lemma 3.10. Let $(r'_p) \in \mathfrak{R}$ be a sequence such that $(r'_p) \leq (r_p)$, $r'_1 = 1$ and the sequence M_p/R'_p satisfies $(M.3)'$ and $(M.4)$. Then for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for every $0 < h < h_0$

$$
\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|} R'_\alpha}{M_\alpha} \|B x^\alpha F[u]\|_{H^s} \leq \varepsilon H_{N-1}^{s,h,(r'_p)}[u].
$$

ii) In the $\{M_p\}$ case, for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for every $0 < h < h_0$

$$
\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{M_\alpha} \|B x^\alpha F[u]\|_{H^s} \leq \varepsilon H_{N-1}^{s,h}[u].
$$

Proof. *i*) Observe that the existence of such sequence $(r'_p) \in \mathfrak{R}$ is given by Lemma 4.2. Let $\alpha \in \mathbb{N}^d$ with $|\alpha| \geq 1$ and let $j = j_\alpha \in \{1, ..., d\}$ such that $\alpha_j > 0$. By Lemma 3.10, there exists $C_1 > 0$ such that

$$\|(B \circ x_j p_\beta(x))(x, D)\|_{\mathcal{L}_b(H^s)} \leq C_1 / R'_\beta.$$

We obtain that $\left\|B\left(x_j p_\beta(x) x^{\alpha - e_j} u^{|\beta|}\right)\right\|_{H^s} \leq \dfrac{C_1}{R'_\beta} \left\|x^{\alpha - e_j} u^{|\beta|}\right\|_{H^s}$. Moreover

$$\left\|x^{\alpha - e_j} u^{|\beta|}\right\|_{H^s} \leq C_s^{|\beta|-1} \left\|x^{\alpha - e_j} u\right\|_{H^s} \|u\|_{H^s}^{|\beta|-1}.$$

Hence

$$\left\|B\left(p_\beta(x) x^\alpha u^{|\beta|}\right)\right\|_{H^s} \leq C_1 \frac{(C_s \|u\|_{H^s})^{|\beta|-1}}{R'_\beta} \left\|x^{\alpha - e_j} u\right\|_{H^s}.$$

Let $C_2 = \displaystyle\sum_{|\beta|=2}^{\infty} \frac{(C_s \|u\|_{H^s})^{|\beta|-1}}{R'_\beta}$. We obtain

$$\begin{aligned} \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B x^\alpha F[u]\|_{H^s} &\leq \sum_{|\alpha|=1}^{N} \sum_{|\beta|=2}^{\infty} \frac{h^{|\alpha|}}{N_\alpha} \left\|B(x^\alpha p_\beta(x) u^{|\beta|})\right\|_{H^s} \\ &\leq C_1 C_2 h \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|-1}}{N_{\alpha - e_j}} \left\|x^{\alpha - e_{j_\alpha}} u\right\|_{H^s} \leq C_3 h H_{N-1}^{s,h,(r'_p)}[u]. \end{aligned}$$

Moreover, for fixed $0 < \varepsilon < 1$, since C_3 does not depend on h, we can find $h_0 = h_0(\varepsilon)$ such that for all $0 < h < h_0$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B x^\alpha F[u]\|_{H^s} \leq \varepsilon H_{N-1}^{s,h,(r'_p)}[u],$$

which complete the proof in the (M_p) case.

ii) In the $\{M_p\}$ case by using Lemma 3.10, one similarly obtains that for every $\tilde{h} > 0$ there exists $C_1 > 0$ such that

$$\left\|B\left(p_\beta(x) x^\alpha u^{|\beta|}\right)\right\|_{H^s} \leq C_1 \left(\tilde{h} C_s \|u\|_{H^s}\right)^{|\beta|-1} \left\|x^{\alpha - e_{j_\alpha}} u\right\|_{H^s}.$$

Fix $\tilde{h}$ such that $\tilde{h} C_s \|u\|_{H^s} < 1/2$. We have

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{M_\alpha} \|B x^\alpha F[u]\|_{H^s} \leq C'_3 h H_{N-1}^{s,h}[u],$$

for a constant C'_3 which is the same for all h. Hence, we obtain the claim in the $\{M_p\}$ case. □

Proof of Theorem 4.3. Fixed $\alpha \in \mathbb{N}^d$ let us multiply both members of (0.1) by x^α. We have $x^\alpha Au = x^\alpha f + x^\alpha F[u]$. Then, introducing commutators we get

$$A(x^\alpha u) = [A, x^\alpha]u + x^\alpha f + x^\alpha F[u]. \tag{4.1}$$

By applying the parametrix B of A to both sides of (4.1) we have

$$x^\alpha u = B[A, x^\alpha]u + B(x^\alpha f) + B(x^\alpha F[u]) + T(x^\alpha u) \tag{4.2}$$

for some *-regularizing operator T. We first consider the (M_p) case. Since $f \in \mathcal{S}^{(M_p)}$, for every $\tilde{h} > 0$ we have $\sup_\alpha \dfrac{\|x^\alpha f\|_{H^s}}{\tilde{h}^{|\alpha|} M_\alpha} < \infty$. Hence, by Lemma 3.4 of [17], there exist $(\tilde{r}_p) \in \mathfrak{R}$ and $C' > 0$ such that $\|x^\alpha f\|_{H^s} \leq C' M_\alpha / \tilde{R}_\alpha$. Obviously, without loss of generality, we can assume that $\tilde{r}_1 = 1$. By Lemma 4.2 we can find $(r'_p) \in \mathfrak{R}$ such that $r'_1 = 1$, $(r'_p) \leq (\tilde{r}_p)$, (r'_p) is smaller than the sequences in Lemmas 3.8 and 3.10, and the sequence $N_p = M_p / R'_p$, for $p \in \mathbb{Z}_+$ and $N_0 = 1$, satisfies $(M.3)'$, $(M.4)$ and $N_1 = 1$. If we multiply (4.2) by $h^{|\alpha|}/N_\alpha$, take Sobolev norms and sum up for $|\alpha| \leq N$, we obtain

$$\begin{aligned} H_N^{s,h,(r'_p)}[u] \;\leq\; & \|u\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B[A, x^\alpha]u\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B(x^\alpha f)\|_{H^s} \\ & + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B(x^\alpha F[u])\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|T(x^\alpha u)\|_{H^s}. \end{aligned}$$

We will estimate each of the terms above. First, since B is bounded on H^s, there exists $C'' > 0$ such that $\|B(x^\alpha f)\|_{H^s} \leq C'' \|x^\alpha f\|_{H^s}$, from what we obtain $\displaystyle\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B(x^\alpha f)\|_{H^s} \leq C''' \sum_{|\alpha|=1}^{\infty} \frac{1}{2^{|\alpha|}} = C_1$ for all $0 < h < 1/2$. To estimate the sum with $T(x^\alpha u)$, since $|\alpha| > 0$ there exists $j = j_\alpha \in \{1, ..., d\}$ such that $\alpha_j \geq 1$. Hence, there exists $C_2 > 0$ such that $\|T \circ x_j\|_{\mathcal{L}_b(H^s)} \leq C_2$. Then we obtain

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|T(x^\alpha u)\|_{H^s} \leq C_2 h \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|-1}}{N_{|\alpha|-1}} \left\|x^{\alpha - e_{j_\alpha}} u\right\|_{H^s} \leq C_3 h H_{N-1}^{s,h,(r'_p)}[u].$$

Since C_3 does not depend on h, for fixed $0 < \varepsilon < 1$ we can find $h_0 = h_0(\varepsilon) < 1/2$ such that for all $0 < h < h_0$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|T(x^\alpha u)\|_{H^s} \leq \varepsilon H_{N-1}^{s,h,(r'_p)}[u].$$

Now, if we use Lemmas 4.6 and 4.5 for fixed $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for all $0 < h < h_0$ we obtain

$$H_N^{s,h,(r'_p)}[u] \leq \|u\|_{H^s} + \varepsilon H_{N-1}^{s,h,(r'_p)}[u] + C_1 + \varepsilon H_{N-1}^{s,h,(r'_p)}[u] + \varepsilon H_{N-1}^{s,h,(r'_p)}[u].$$

By iterating this estimate and possibly shrinking ε we obtain that $\sum_{|\alpha|=0}^{\infty} \frac{h^{|\alpha|}}{N_\alpha}\|x^\alpha u\|_{H^s}$ is finite for some sufficiently small h. If $\tilde{h} > 0$ is arbitrary but fixed, there exists $\tilde{C} > 0$ such that $\tilde{h}^p \leq \tilde{C} h^p R'_p$, for all $p \in \mathbb{Z}_+$. Hence the sum $\sum_{|\alpha|=0}^{\infty} \frac{\tilde{h}^{|\alpha|}}{M_\alpha}\|x^\alpha u\|_{H^s}$ converges. This completes the proof in the (M_p) case. The $\{M_p\}$ case is completely similar. We leave the details to the reader. □

Now we prove Theorem 4.4.

Lemma 4.7. *Let $A = a(x, D)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) operator and let B be its parametrix. Then the following properties hold:*

i) In the (M_p) case, let $(r_p) \in \mathfrak{R}$ be the sequence in Lemma 3.8. Let $(r'_p) \in \mathfrak{R}$ be a sequence such that $(r'_p) \leq (r_p)$, $r'_1 = 1$ and the sequence M_p/R'_p satisfies $(M.3)'$ and $(M.4)$. Then for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for every $0 < h < h_0$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|} R'_\alpha}{M_\alpha} \|B[A, \partial^\alpha]u\|_{H^s} \leq \varepsilon E_{N-1}^{s,h,(r'_p)}[u].$$

ii) In the $\{M_p\}$ case, for each $0 < \varepsilon < 1$ there exists $h_0 = h_0(\varepsilon)$ such that for all $0 < h < h_0(\varepsilon)$

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{M_\alpha} \|B[A, \partial^\alpha]u\|_{H^s} \leq \varepsilon E_{N-1}^{s,h}[u].$$

Proof. First we prove the (M_p) case. As before, put $N_p = M_p/R'_p$, for $p \in \mathbb{Z}_+$ and $N_0 = 1$ (the existence of such $(r'_p) \in \mathfrak{R}$ is given by Lemma 4.2). Observe that

$$B[A, \partial^\alpha]u = -\sum_{\substack{\beta \leq \alpha \\ \beta \neq 0}} \binom{\alpha}{\beta} B(\partial_x^\beta a)(x, D)\partial_x^{\alpha-\beta} u.$$

By Lemma 3.8, there exists $C > 0$ such that $\left\|B(\partial_x^\beta a)(x, D)\right\|_{\mathcal{L}_b(H^s)} \leq C N_\beta$. Let $0 < \varepsilon < 1$ be fixed. Choose $0 < h_0 < 1/2$ such that $h_0 < \varepsilon \left(2C \sum_{|\beta|=1}^{\infty} 2^{-|\beta|+1}\right)^{-1}$. For $0 < h < h_0$ we obtain

$$\begin{aligned}
&\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \|B[A, \partial^\alpha]u\|_{H^s} \\
&\leq \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \sum_{\substack{\beta \leq \alpha \\ \beta \neq 0}} \binom{\alpha}{\beta} \left\|B(\partial_x^\beta a)(x, D)\partial_x^{\alpha-\beta} u\right\|_{H^s}
\end{aligned}$$

$$\leq C\sum_{|\alpha|=1}^{N}\sum_{\substack{\beta\leq\alpha\\ \beta\neq 0}}\frac{h^{|\alpha|}N_\beta}{N_\alpha}\binom{\alpha}{\beta}\left\|\partial_x^{\alpha-\beta}u\right\|_{H^s}\leq C\sum_{|\beta|=1}^{N}h^{|\beta|}\sum_{\substack{\alpha\geq\beta\\ |\alpha|\leq N}}\frac{h^{|\alpha|-|\beta|}}{N_{\alpha-\beta}}\left\|\partial_x^{\alpha-\beta}u\right\|_{H^s}$$

$$\leq \varepsilon E_{N-1}^{s,h,(r_p')}[u],$$

where in the third inequality, we used $(M.4)$ for N_p and the fact $\binom{\alpha}{\beta}\leq\binom{|\alpha|}{|\beta|}$. This completes the proof in the (M_p) case. For the $\{M_p\}$ case, let $\varepsilon>0$. By Lemma 3.8, there exist $h_1, C>0$ such that $\left\|B(\partial_x^\beta a)(x,D)\right\|_{\mathcal{L}_b(H^s)}\leq Ch_1^{|\beta|}M_\beta$. Choose $h_0>0$ such that $h_0h_1<1/2$ and $h_0h_1\leq\varepsilon\left(2C\sum_{|\beta|=1}^{\infty}2^{-|\beta|+1}\right)^{-1}$. Then, for $0<h<h_0$, similarly as before, we obtain

$$\begin{aligned}\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{M_\alpha}\left\|B[A,\partial^\alpha]u\right\|_{H^s} &\leq C\sum_{|\beta|=1}^{N}(hh_1)^{|\beta|}\sum_{\substack{\alpha\geq\beta\\ |\alpha|\leq N}}\frac{h^{|\alpha|-|\beta|}}{M_{\alpha-\beta}}\left\|\partial_x^{\alpha-\beta}u\right\|_{H^s}\\ &\leq \varepsilon E_{N-1}^{s,h}[u],\end{aligned}$$

which completes the proof. □

Lemma 4.8. *Let $A=a(x,D)$ be $(\tilde{M}_p)$-elliptic (resp. $\{\tilde{M}_p\}$-elliptic) operator and let B be its parametrix. Let $F[u]=p(x)u^l$, for some $l\geq 2$, $l\in\mathbb{N}$. Then the following properties hold:*

i) In the (M_p) case, let $(r_p)\in\mathfrak{R}$ be the sequence in Lemma 3.10. Let $(r_p')\in\mathfrak{R}$ be a sequence such that $(r_p')\leq(r_p)$, $r_1'=1$ and the sequence M_p/R_p' satisfies $(M.3)'$ and $(M.4)$. Then for each $0<\varepsilon<1$ there exists $h_0=h_0(\varepsilon)$ such that for every $0<h<h_0$

$$\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}R_\alpha'}{M_\alpha}\left\|B\left(\partial^\alpha(p(x)u^l)\right)\right\|_{H^s}\leq\varepsilon\left(E_{N-1}^{s,h,(r_p')}[u]\right)^l.$$

ii) In the $\{M_p\}$ case, for each $0<\varepsilon<1$ there exists $h_0=h_0(\varepsilon)$ such that for every $0<h<h_0$

$$\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{M_\alpha}\left\|B\left(\partial^\alpha(p(x)u^l)\right)\right\|_{H^s}\leq\varepsilon\left(E_{N-1}^{s,h}[u]\right)^l.$$

Proof. Observe that

$$B\left(\partial^\alpha(p(x)u^l)\right)=B\left(p(x)\partial^\alpha u^l\right)+\sum_{\substack{\gamma\leq\alpha\\ \gamma\neq 0}}\binom{\alpha}{\gamma}B\left(\partial^\gamma p(x)\partial^{\alpha-\gamma}u^l\right)$$

First we consider the (M_p) case. As before, put $N_p=M_p/R_p'$, for $p\in\mathbb{Z}_+$ and $N_0=1$. Since $|\alpha|\geq 1$, there exists $j=j_\alpha\in\{1,...,d\}$ such that $\alpha_j>0$. By Lemma 3.10, there exists $C_1>0$ such that $\left\|\left(B\circ p(x)\partial_j\right)(x,D)\right\|_{\mathcal{L}_b(H^s)}\leq C_1$. Then we have

$$\left\|B\left(p(x)\partial^\alpha u^l\right)\right\|_{H^s}\leq C_1\left\|\partial^{\alpha-e_j}u^l\right\|_{H^s}$$

$$\leq C_2 \sum_{\nu^{(1)}+\ldots+\nu^{(l)}=\alpha-e_j} \frac{(\alpha-e_j)!}{\nu^{(1)}!\cdot\ldots\cdot\nu^{(l)}!} \prod_{k=1}^{l} \left\|\partial^{\nu^{(k)}} u\right\|_{H^s}.$$

Observe that, by $(M.4)$,

$$\frac{(\alpha-e_j)!}{\nu^{(1)}!\cdot\ldots\cdot\nu^{(l)}!}\cdot\frac{h^{|\alpha|}}{N_\alpha} \leq \frac{hN_{|\alpha|-1}}{N_\alpha}\prod_{k=1}^{l}\frac{h^{|\nu^{(k)}|}}{N_{\nu^{(k)}}} \leq h\prod_{k=1}^{l}\frac{h^{|\nu^{(k)}|}}{N_{\nu^{(k)}}}.$$

We obtain

$$\begin{aligned}\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{N_\alpha}\left\|B\left(p(x)\partial^\alpha u^l\right)\right\|_{H^s} &\leq C_2 h\sum_{|\alpha|=1}^{N}\sum_{\nu^{(1)}+\ldots+\nu^{(l)}=\alpha-e_{j_\alpha}}\prod_{k=1}^{l}\frac{h^{|\nu^{(k)}|}}{N_{\nu^{(k)}}}\left\|\partial^{\nu^{(k)}}u\right\|_{H^s}\\ &\leq dC_2h\left(E_{N-1}^{s,h,(r'_p)}[u]\right)^l.\end{aligned}$$

Since C_2 does not depend on h, for fixed $0<\varepsilon<1$ we can take $h_0=\varepsilon/(dC_2)$. Then for all $h<h_0$ we obtain $\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{N_\alpha}\left\|B\left(p(x)\partial^\alpha u^l\right)\right\|_{H^s}\leq\varepsilon\left(E_{N-1}^{s,h,(r'_p)}[u]\right)^l$. One easily verifies that the functions $p'_\beta(x)=\partial_x^\beta p(x)$, $\beta\in\mathbb{N}^d$ satisfy (1.3). Lemma 3.10 implies that there exists $C_1>0$ such that $\|B\circ\partial^\gamma p(x)\|_{\mathcal{L}_b(H^s)}\leq C_1/R'_\gamma$. For $h<1$ we obtain

$$\begin{aligned}&\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{N_\alpha}\sum_{\substack{\gamma\leq\alpha\\ \gamma\neq 0}}\binom{\alpha}{\gamma}\left\|B\left(\partial^\gamma p(x)\partial^{\alpha-\gamma}u^l\right)\right\|_{H^s}\\ &\leq C_1h\sum_{|\alpha|=1}^{N}\sum_{\substack{\gamma\leq\alpha\\ \gamma\neq 0}}\binom{\alpha}{\gamma}\frac{h^{|\alpha|-1}}{N_\alpha R'_\gamma}\left\|\partial^{\alpha-\gamma}u^l\right\|_{H^s}\\ &\leq C'_2h\sum_{|\alpha|=1}^{N}\sum_{\substack{\gamma\leq\alpha\\ \gamma\neq 0}}\binom{\alpha}{\gamma}\frac{h^{|\alpha|-1}}{N_\alpha R'_\gamma}\sum_{\nu^{(1)}+\ldots+\nu^{(l)}=\alpha-\gamma}\frac{(\alpha-\gamma)!}{\nu^{(1)}!\cdot\ldots\cdot\nu^{(l)}!}\prod_{k=1}^{l}\left\|\partial^{\nu^{(k)}}u\right\|_{H^s}\\ &\leq C'_2h\sum_{|\gamma|=1}^{N}\frac{1}{R'_\gamma}\sum_{\substack{\alpha\geq\gamma\\ |\alpha|\leq N}}\frac{h^{|\alpha|-1}}{N_\alpha}\sum_{\nu^{(1)}+\ldots+\nu^{(l)}=\alpha-\gamma}\frac{\alpha!}{\nu^{(1)}!\cdot\ldots\cdot\nu^{(l)}!\gamma!}\prod_{k=1}^{l}\left\|\partial^{\nu^{(k)}}u\right\|_{H^s}\\ &\leq C'_2h\sum_{|\gamma|=1}^{N}\frac{1}{N_\gamma R'_\gamma}\sum_{\substack{\alpha\geq\gamma\\ |\alpha|\leq N}}\sum_{\nu^{(1)}+\ldots+\nu^{(l)}=\alpha-\gamma}\prod_{k=1}^{l}\frac{h^{|\nu^{(k)}|}}{N_{\nu^{(k)}}}\left\|\partial^{\nu^{(k)}}u\right\|_{H^s}\\ &\leq C'_2h\sum_{|\gamma|=1}^{N}\frac{1}{M_\gamma}\sum_{|\nu^{(1)}|+\ldots+|\nu^{(l)}|=0}^{N-1}\prod_{k=1}^{l}\frac{h^{|\nu^{(k)}|}}{N_{\nu^{(k)}}}\left\|\partial^{\nu^{(k)}}u\right\|_{H^s}\leq C'_3h\left(E_{N-1}^{s,h,(r'_p)}[u]\right)^l.\end{aligned}$$

Since C'_3 does not depend on h, for fixed $0<\varepsilon<1$ we can choose $h_0=\varepsilon/C'_3$. Then,

for all $0 < h < h_0$, we have

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \sum_{\substack{\gamma\leq\alpha \\ |\gamma|\neq 0}} \binom{\alpha}{\gamma} \left\| B\left(\partial^\gamma p(x)\partial^{\alpha-\gamma}u^l\right)\right\|_{H^s} \leq \varepsilon \left(E_{N-1}^{s,h,(r'_p)}[u]\right)^l,$$

which, combined with the above estimate for $\displaystyle\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha} \left\| B\left(p(x)\partial^\alpha u^l\right)\right\|_{H^s}$, completes the proof in the (M_p) case by shrinking h_0 if necessary. In the $\{M_p\}$ case the proof is similar. □

Proof of Theorem 4.4. i) When $F[u]$ is a finite sum it is clear that it is enough to prove the theorem when $F[u] = p(x)u^l$, $l \geq 2$, $l \in \mathbb{N}$. Differentiating both terms of (0.1), we have $\partial^\alpha Au = \partial^\alpha f + \partial^\alpha F[u]$, from which we obtain

$$A(\partial^\alpha u) = [A, \partial^\alpha]u + \partial^\alpha f + \partial^\alpha F[u].$$

Hence, we have

$$\partial^\alpha u = B[A, \partial^\alpha]u + B(\partial^\alpha f) + B(\partial^\alpha F[u]) + T(\partial^\alpha u). \tag{4.3}$$

We consider the (M_p) case. Since $f \in \mathcal{S}^{(M_p)}$, for every $\tilde{h} > 0$, $\displaystyle\sup_\alpha \frac{\|\partial^\alpha f\|_{H^s}}{\tilde{h}^{|\alpha|}M_\alpha}$ is bounded. Hence, by Lemma 3.4 of [17], there exist $(\tilde{r}_p) \in \mathfrak{R}$ and $C' > 0$ such that $\|\partial^\alpha f\|_{H^s} \leq C'M_\alpha/\tilde{R}_\alpha$. Obviously, without losing generality, we can assume that $\tilde{r}_1 = 1$. By Lemma 4.2 we can find $(r'_p) \in \mathfrak{R}$ such that $r'_1 = 1$, $(r'_p) \leq (\tilde{r}_p)$, (r'_p) is smaller than the sequences in Lemmas 3.8 and 3.10 and the sequence $N_p = M_p/R'_p$, for $p \in \mathbb{Z}_+$ and $N_0 = 1$, satisfies $(M.3)'$, $(M.4)$ and $N_1 = 1$. If we multiply (4.3) by $h^{|\alpha|}/N_\alpha$, take Sobolev norms and sum up for $|\alpha| \leq N$, we obtain

$$\begin{aligned} E_N^{s,h,(r'_p)}[u] \quad \leq \quad & \|u\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|B[A, \partial^\alpha]u\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|B(\partial^\alpha f)\|_{H^s} \\ & + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|B(\partial^\alpha F[u])\|_{H^s} + \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|T(\partial^\alpha u)\|_{H^s}. \end{aligned}$$

We estimate each of the terms above. By the growth estimate for the symbol of B (3.2) there exists $C'' > 0$ such that $\|B(\partial^\alpha f)\|_{H^s} \leq C''\|\partial^\alpha f\|_{H^s}$. Hence $\displaystyle\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|B(\partial^\alpha f)\|_{H^s} \leq C''' \sum_{|\alpha|=1}^{\infty} \frac{1}{2^{|\alpha|}} = C_1$ for all $0 < h < 1/2$. To estimate the sum with $T(\partial^\alpha u)$, since $|\alpha| > 0$ there exists $j_\alpha \in \{1, ..., d\}$ such that $\alpha_{j_\alpha} \geq 1$. Hence there exists $C_2 > 0$ such that $\|T \circ \partial^{e_{j_\alpha}}\|_{\mathcal{L}_b(H^s)} \leq C_2$. Then we obtain

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{N_\alpha}\|T(\partial^\alpha u)\|_{H^s} \leq C_2 h \sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|-1}}{N_{|\alpha|-1}} \left\|\partial^{\alpha-e_{j_\alpha}}u\right\|_{H^s} \leq C_3 h E_{N-1}^{s,h,(r'_p)}[u].$$

Since C_3 does not depend on h, for fixed $0<\varepsilon<1$ we can find $h_0=h_0(\varepsilon)<1/2$ such that for all $0<h<h_0$

$$\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{N_\alpha}\|T(\partial^\alpha u)\|_{H^s}\leq \varepsilon E_{N-1}^{s,h,(r'_p)}[u].$$

For fixed $0<\varepsilon<1$, by Lemmas 4.7 and 4.8 for the chosen (r'_p), we can find $h_0=h_0(\varepsilon)<1/2$ such that for all $0<h<h_0$, we have

$$E_N^{s,h,(r'_p)}[u] \;\leq\; \|u\|_{H^s}+\varepsilon E_{N-1}^{s,h,(r'_p)}[u]+C_1+\varepsilon\left(E_{N-1}^{s,h,(r'_p)}[u]\right)^l+\varepsilon E_{N-1}^{s,h,(r'_p)}[u].$$

By iterating this estimate one obtains that $\|\partial^\alpha u\|_{H^s}$, $\alpha\in\mathbb{N}^d$, are finite and by shrinking ε if necessary, that the sum $\displaystyle\sum_{|\alpha|=0}^{\infty}\frac{h^{|\alpha|}}{N_\alpha}\|\partial^\alpha u\|_{H^s}$ converges, for some, small enough, h. If $\tilde h>0$ is arbitrary but fixed there exists $\tilde C>0$ such that $\tilde h^p\leq Ch^pR'_p$ for all $p\in\mathbb{Z}_+$. Hence $\displaystyle\sum_{|\alpha|=0}^{\infty}\frac{\tilde h^{|\alpha|}}{M_\alpha}\|\partial^\alpha u\|_{H^s}$ converges. This completes the proof in the (M_p) case. The proof in the $\{M_p\}$ case is similar and we omit it.

To prove $ii)$ we consider first the (M_p) case. Proceed as in the proof $i)$ to obtain

$$\begin{aligned}\sum_{|\alpha|\leq N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|\partial^\alpha u\|_{H^s} \;\leq\; & \|u\|_{H^s}+\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|B[A,\partial^\alpha]u\|_{H^s}+\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|B(\partial^\alpha f)\|_{H^s}\\ &+\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|B(\partial^\alpha F[u])\|_{H^s}+\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|T(\partial^\alpha u)\|_{H^s}.\end{aligned}$$

By Lemma 3.9, there exists $l\geq 1$ such that for each $\tilde h>0$ there exists $C_1>0$ such that $\|(B\circ D^\alpha p_\beta(x)\partial^\gamma)(x,D)\|_{\mathcal{L}(H^s)}\leq C_1\tilde h^{|\beta|}l^{|\gamma|}\tilde M_\gamma$. For $0<h<1/(4l)$, we have

$$\begin{aligned}\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\|B(\partial^\alpha F[u])\|_{H^s} \;&\leq\; \sum_{|\beta|=2}^{\infty}\sum_{|\alpha|=1}^{N}\sum_{\gamma\leq\alpha}\binom{\alpha}{\gamma}\frac{h^{|\alpha|}}{\tilde M_\alpha}\left\|B(\partial^\gamma p_\beta\partial^{\alpha-\gamma}u^{|\beta|})\right\|_{H^s}\\ &\leq\; C_1\sum_{|\beta|=2}^{\infty}\sum_{|\alpha|=1}^{N}\frac{\tilde h^{|\beta|}\left\|u^{|\beta|}\right\|_{H^s}}{2^{|\alpha|}}\leq C_2,\end{aligned}$$

where in the last inequality we used that $\left\|u^{|\beta|}\right\|_{H^s}\leq C_s^{|\beta|-1}\|u\|_{H^s}^{|\beta|}$ and chose $\tilde h\leq 1/(2C_s\|u\|_{H^s})$. The sequence $\tilde M_p$ satisfies $(M.4)$, so by analogous technique as in the proof of Lemma 4.7 one can prove that for each $0<\varepsilon<1$ there exists $h_0=h_0(\varepsilon)<1/2$ such that for every $0<h<h_0$

$$\sum_{|\alpha|=1}^{N}\frac{h^{|\alpha|}}{\tilde M_\alpha}\left\|B[A,\partial^\alpha]u\right\|_{H^s}\leq\varepsilon\sum_{|\alpha|\leq N-1}\frac{h^{|\alpha|}}{\tilde M_\alpha}\left\|\partial^\alpha u\right\|_{H^s}.$$

Also, similarly as in the proof of i), we have that for $0 < h < h_0(\varepsilon)$,

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|B(\partial^\alpha f)\|_{H^s} \leq C_3$$

and

$$\sum_{|\alpha|=1}^{N} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|T(\partial^\alpha u)\|_{H^s} \leq \varepsilon \sum_{|\alpha|\leq N-1} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|\partial^\alpha u\|_{H^s}.$$

Hence, for $0 < h < h_0(\varepsilon)$, for sufficiently small $h_0(\varepsilon)$, we have

$$\sum_{|\alpha|\leq N} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|\partial^\alpha u\|_{H^s} \leq \|u\|_{H^s} + C_4 + 2\varepsilon \sum_{|\alpha|\leq N-1} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|\partial^\alpha u\|_{H^s}.$$

By iterating this estimate and possibly shrinking ε we obtain that $\sum_{|\alpha|=0}^{\infty} \frac{h^{|\alpha|}}{\tilde{M}_\alpha} \|\partial^\alpha u\|_{H^s}$ is finite for some sufficiently small h, which finishes the proof in the (M_p) case. The $\{M_p\}$ case is completely analogous. □

References

[1] H. A. Biagioni and T. Gramchev, *Fractional derivative estimates in Gevrey classes, global regularity and decay for solutions to semilinear equations in* $\mathbb{R}^n$, J. Differential Equations, **194** (2003), 140–165.

[2] M. Cappiello, *Gelfand spaces and pseudodifferential operators of infinite order in* $\mathbb{R}^n$, Ann. Univ Ferrara, Sez. VII, Sc. Mat., **48** (2002), 75-97.

[3] M. Cappiello, *Pseudodifferential parametrices of infinite order for SG-hyperbolic problems*, Rend. Sem. Mat. Univ. Pol. Torino, **61** 4 (2003), 411-441.

[4] M. Cappiello, T. Gramchev and L. Rodino, *Super-exponential decay and holomorphic extensions for semilinear equations with polynomial coefficients*, J. Funct. Anal., **237** (2006), 634–654.

[5] M. Cappiello, T. Gramchev and L. Rodino, *Sub-exponential decay and uniform holomorphic extensions for semilinear pseudodifferential equations*, Comm. Partial Differential Equations, **35** (2010) n. 5, 846–877.

[6] M. Cappiello, T. Gramchev and L. Rodino, *Entire extensions and exponential decay for semilinear elliptic equations*, J. Anal. Math., **111** (2010), 339–367.

[7] M. Cappiello and F. Nicola, *Regularity and decay of solutions of nonlinear harmonic oscillators*, Adv. Math. **229** (2012), 1266–1299.

[8] M. Cappiello, S. Pilipović and B. Prangoski, *Parametrices and hypoellipticity for pseudodifferential operators on spaces of tempered ultradistributions*, J. Pseudo-Differ. Oper. Appl., online July 2014, http://dx.doi.org/10.1007/s11868-014-0095-3

[9] J. Chung, S.Y. Chung and D. Kim, *Characterizations of the Gelfand-Shilov spaces via Fourier transforms*, Proc. Am. Math. Soc. **124** 7 (1996), 2101-2108

[10] G. M. Constantine and T. H. Savits, *A multivariate Faà di Bruno formula with applications*, Trans. Amer. Math. Soc., **348** 2 (1996), 503-520.

[11] T. Gramchev, *Perturbative methods in scales of Banach spaces: applications for Gevrey regularity of solutions to semilinear partial differential equations*, Rend. Sem. Mat. Univ. Pol. Torino, **61** (2003) no. 2, 101–134.

[12] R. Carmichael, A. Kamiński and S. Pilipović, *Boundary Values and Convolution in Ultradistribution Spaces*, World Scientific Publishing Co. Pte. Ltd., 2007

[13] I.M. Gelfand and G.E. Shilov, *Generalized functions II*, Academic Press, New York, 1968.

[14] H. Komatsu, *Ultradistributions, I: Structure theorems and a characterization*, J. Fac. Sci. Univ. Tokyo, Sect. IA Math., **20** 1 (1973), 25-105.

[15] H. Komatsu, *Ultradistributions, II: The kernel theorem and ultradistributions with support in submanifold*, J. Fac. Sci. Univ. Tokyo, Sect. IA Math., **24** 3 (1977), 607-628.

[16] H. Komatsu, *The implicit function theorem for ultradifferentiable mappings*, Proc. Japan Acad., Ser. A **55** 3 (1979), 69-72.

[17] H. Komatsu, *Ultradistributions, III: Vector valued ultradistributions and the theory of kernels*, J. Fac. Sci. Univ. Tokyo, Sect. IA Math., **29** 3 (1982), 653-717.

[18] F. Nicola and L. Rodino, *Global Psedo-Differential Calculus on Euclidean Spaces*, Vol. 4. Birkhäuser Basel, 2010

[19] S. Pilipovic, *Tempered ultradistributions*, Boll. Un. Mat. Ital. B (7) **2** (1988), no. 2, 235-251.

[20] S. Pilipović, *Characterizations of bounded sets in spaces of ultradistributions*, Proc. Amer. Math. Soc., **120** (1994) no. 4, 1191–1206.

[21] S. Pilipović and B. Prangoski, *Anti-Wick and Weyl quantization on ultradistribution spaces*, J. Math. Pures Appl., online April 2014, http://dx.doi.org/10.1016/j.matpur.2014.04.011

[22] B. Prangoski, *Laplace transform in spaces of ultradistributions*, Filomat **27** 5 (2013), 747-760.

[23] B. Prangoski, *Pseudodifferential operators of infinite order in spaces of tempered ultradistributions*, J. Pseudo-Differ. Oper. Appl. **4** (2013), 495-549.